

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT

(10) International Publication Number
WO 01/90122 A2

(51) International Patent Classification: C07H

(74) Agents: HENDERSON, Melodie, W. et al.: Genaisance Pharmaceuticals, Inc., Five Science Park, New Haven, CT 06511 (US).

(21) International Application Number: PCT/US01/16905

(22) International Filing Date: 23 May 2001 (23.05.2001)

(81) Designated States (national): AE, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 60/206,353 23 May 2000 (23.05.2000) US

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): GENAISANCE PHARMACEUTICALS, INC. (US/US); Five Science Park, New Haven, CT 06511 (US).

Published:

— without international search report and to be republished upon receipt of that report

(72) Inventors; and
(75) Inventors/Applicants (for US only): CHEW, Anne (US/US); 1477 Beacon Street, #64, Brookline, MA 02446 (US). CHOI, Julie, Y. (US/US); 38 Elizabeth Street, West Haven, CT 06516 (US). MESSER, Chad (US/US); 41 Manor Street, Hamden, CT 06517 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/90122 A2

(54) Title: HAPLOTYPES OF THE CTLA4 GENE

(57) Abstract: Novel genetic variants of the Cytotoxic T-Lymphocyte-Associated Protein 4 (CTLA4) gene are described. Various genotypes, haplotypes, and haplotype pairs that exist in the general United States population are disclosed for the CTLA4 gene. Compositions and methods for haplotyping and/or genotyping the CTLA4 gene in an individual are also disclosed. Polynucleotides defined by the sequence the haplotypes disclosed herein are also described.

HAPLOTYPES OF THE CTLA4 GENE

RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application Serial No. 60/206,353 filed May 23, 2000.

FIELD OF THE INVENTION

This invention relates to variation in genes that encode pharmaceutically-important proteins. In particular, this invention provides genetic variants of the human cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene and methods for identifying which variant(s) of this gene is/are possessed by an individual.

BACKGROUND OF THE INVENTION

Current methods for identifying pharmaceuticals to treat disease often start by identifying, cloning, and expressing an important target protein related to the disease. A determination of whether an agonist or antagonist is needed to produce an effect that may benefit a patient with the disease is then made. Then, vast numbers of compounds are screened against the target protein to find new potential drugs. The desired outcome of this process is a lead compound that is specific for the target, thereby reducing the incidence of the undesired side effects usually caused by activity at non-intended targets. The lead compound identified in this screening process then undergoes further *in vitro* and *in vivo* testing to determine its absorption, disposition, metabolism and toxicological profiles. Typically, this testing involves use of cell lines and animal models with limited, if any, genetic diversity.

What this approach fails to consider, however, is that natural genetic variability exists between individuals in any and every population with respect to pharmaceutically-important proteins, including the protein targets of candidate drugs, the enzymes that metabolize these drugs and the proteins whose activity is modulated by such drug targets. Subtle alteration(s) in the primary nucleotide sequence of a gene encoding a pharmaceutically-important protein may be manifested as significant variation in expression, structure and/or function of the protein. Such alterations may explain the relatively high degree of uncertainty inherent in the treatment of individuals with a drug whose design is based upon a single representative example of the target or enzyme(s) involved in metabolizing the drug. For example, it is well-established that some drugs frequently have lower efficacy in some individuals than others, which means such individuals and their physicians must weigh the possible benefit of a larger dosage against a greater risk of side effects. Also, there is significant variation in how well people metabolize drugs and other exogenous chemicals, resulting in substantial interindividual variation in the toxicity and/or efficacy of such exogenous substances (Evans et al., 1999, *Science* 286:487-491). This variability in efficacy or toxicity of a drug in genetically-diverse patients makes many drugs ineffective or even dangerous in certain groups of the population, leading to the failure of such drugs in clinical trials or their early withdrawal from the market even though they could be highly beneficial for

myocardial failure caused by T-cell blast infiltration (Waterhouse et al., *Science* 1995; 270:985-988). These results suggest that CTLA4 may play an inhibitory role in regulating lymphocyte expansion.

The cytotoxic T-lymphocyte-associated protein 4 gene is located on chromosome 2q33 and contains 4 exons that encode a 223 amino acid protein. Reference sequences for the CTLA4 gene (Genaissance Reference No. 743670; SEQ ID NO: 1), coding sequence (GenBank Accession No:NM_005214.1), and protein are shown in Figures 1, 2 and 3, respectively.

Several variations have been identified in the CTLA4 gene that may be related to various disorders. A polymorphism of adenine or guanine at nucleotide position 37902 in Figure 1 results in an amino acid variation of threonine or alanine at amino acid position 17 in Figure 3 (HGBASE:SNP000000387). Donner et al., (*J Clin Endocrinol. Metab* 1997; 82:4130-4132) reported that patients with Hashimoto thyroiditis had higher frequencies of the Thr17Ala mutation. However, there was no significant variation in patients with Addison's Disease and control subjects. The Thr17Ala mutation has also been shown to be associated with Grave's Disease in a dataset of white Caucasian subjects (Heward et al., *J Clin Endocrinol. Metab* 1999; 84:2398-2401). Djilali-Saiah et al. (*Gut* 1998; 43:187-189) found that in French Caucasian patients with Celiac disease, which is characterized by immunologically mediated intestinal injury following ingestion of gluten, the Thr17Ala mutation was found with greater frequency in patients than in controls. These results suggest that the location of this polymorphism on the gene is critical to the function of the CTLA4 protein.

The association between Thr17Ala polymorphism and insulin dependent diabetes mellitus (IDDM) has also been studied in numerous populations. Associations have been observed several populations including Asian, Mexican-American, and certain Caucasian populations (Awata et al., *Diabetes* 1998; 47:128-129; Donner et al., *J Clin Endocrinol. Metab* 1997; 82:143-146; Lee et al., *Clin Endocrinol. (Oxf)* 2000; 52:153-157; Marron et al., *Hum Mol. Genet* 1997; 6:1275-1282), while a lack of association has been observed in other Caucasian groups (Owerbach et al., *Diabetes* 1997; 46:1069-1074). Chistiakov et al. tested whether or not the CTLA4 gene was a susceptibility marker for IDDM in a Russian population. In the case of the codon 17 polymorphism, the alanine allele was associated with the disease. The authors also examined an (AT)_n microsatellite marker in the 3' untranslated region of the gene, and examined the transmission of 17 alleles varying from 92 to 130 bp in length. The transmission of three alleles was significantly different for diabetic and non-diabetic offspring (Chistiakov et al., *supra*). Therefore, the CTLA4 gene is associated with IDDM in a Russian population.

Because of the potential for variation in the CTLA4 gene to affect the expression and function of the encoded protein, it would be useful to know whether additional polymorphisms exist in the CTLA4 gene, as well as how such polymorphisms are combined in different copies of the gene. Such information could be applied for studying the biological function of CTLA4 as well as in identifying drugs targeting this protein for the treatment of disorders related to its abnormal expression or function.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates a reference sequence for the CTLA4 gene (Genaissance Reference No. 743670; contiguous lines; SEQ ID NO:1), with the start and stop positions of each region of coding sequence indicated with a bracket ([or]) and the numerical position below the sequence and the polymorphic site(s) and polymorphism(s) identified by Applicants in a reference population indicated by the variant nucleotide positioned below the polymorphic site in the sequence. SEQ ID NO:36 is equivalent to Figure 1, with the two alternative allelic variants of each polymorphic site indicated by the appropriate nucleotide symbol (R= G or A, Y= T or C, M= A or C, K= G or T, S= G or C, and W= A or T; WIPO standard ST.25).

Figure 2 illustrates a reference sequence for the CTLA4 coding sequence (contiguous lines; SEQ ID NO:2), with the polymorphic site(s) and polymorphism(s) identified by Applicants in a reference population indicated by the variant nucleotide positioned below the polymorphic site in the sequence.

Figure 3 illustrates a reference sequence for the CTLA4 protein (contiguous lines; SEQ ID NO:3), with the variant amino acid(s) caused by the polymorphism(s) of Figure 2 positioned below the polymorphic site in the sequence.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is based on the discovery of novel variants of the CTLA4 gene. As described in more detail below, the inventors herein discovered 8 isogenes of the CTLA4 gene by characterizing the CTLA4 gene found in genomic DNAs isolated from an Index Repository that contains immortalized cell lines from one chimpanzee and 93 human individuals. The human individuals included a reference population of 79 unrelated individuals self-identified as belonging to one of four major population groups: Caucasian (21 individuals), African descent (20 individuals), Asian (20 individuals), or Hispanic/Latino (18 individuals). To the extent possible, the members of this reference population were organized into population subgroups by their self-identified ethnogeographic origin as shown in Table 1 below.

Gene - A segment of DNA that contains all the information for the regulated biosynthesis of an RNA product, including promoters, exons, introns, and other untranslated regions that control expression.

Genotype - An unphased 5' to 3' sequence of nucleotide pair(s) found at one or more polymorphic sites in a locus on a pair of homologous chromosomes in an individual. As used herein, genotype includes a full-genotype and/or a sub-genotype as described below.

Full-genotype - The unphased 5' to 3' sequence of nucleotide pairs found at all polymorphic sites examined herein in a locus on a pair of homologous chromosomes in a single individual.

Sub-genotype - The unphased 5' to 3' sequence of nucleotides seen at a subset of the polymorphic sites examined herein in a locus on a pair of homologous chromosomes in a single individual.

Genotyping - A process for determining a genotype of an individual.

Haplotype - A 5' to 3' sequence of nucleotides found at one or more polymorphic sites in a locus on a single chromosome from a single individual. As used herein, haplotype includes a full-haplotype and/or a sub-haplotype as described below.

Full-haplotype - The 5' to 3' sequence of nucleotides found at all polymorphic sites examined herein in a locus on a single chromosome from a single individual.

Sub-haplotype - The 5' to 3' sequence of nucleotides seen at a subset of the polymorphic sites examined herein in a locus on a single chromosome from a single individual.

Haplotype pair - The two haplotypes found for a locus in a single individual.

Haplotyping - A process for determining one or more haplotypes in an individual and includes use of family pedigrees, molecular techniques and/or statistical inference.

Haplotype data - Information concerning one or more of the following for a specific gene: a listing of the haplotype pairs in each individual in a population; a listing of the different haplotypes in a population; frequency of each haplotype in that or other populations, and any known associations between one or more haplotypes and a trait.

Isoform - A particular form of a gene, mRNA, cDNA or the protein encoded thereby, distinguished from other forms by its particular sequence and/or structure.

Isogene - One of the isoforms of a gene found in a population. An isogene contains all of the polymorphisms present in the particular isoform of the gene.

Isolated - As applied to a biological molecule such as RNA, DNA, oligonucleotide, or protein, isolated means the molecule is substantially free of other biological molecules such as nucleic acids, proteins, lipids, carbohydrates, or other material such as cellular debris and growth media. Generally, the term "isolated" is not intended to refer to a complete absence of such material or to absence of water, buffers, or salts, unless they are present in amounts that substantially interfere with the methods of the present invention.

Locus - A location on a chromosome or DNA molecule corresponding to a gene or a physical or phenotypic feature.

Naturally-occurring – A term used to designate that the object it is applied to, e.g., naturally-occurring polynucleotide or polypeptide, can be isolated from a source in nature and which has not been intentionally modified by man.

Nucleotide pair – The nucleotides found at a polymorphic site on the two copies of a chromosome from an individual.

Phased – As applied to a sequence of nucleotide pairs for two or more polymorphic sites in a locus, phased means the combination of nucleotides present at those polymorphic sites on a single copy of the locus is known.

Polymorphic site (PS) – A position within a locus at which at least two alternative sequences are found in a population, the most frequent of which has a frequency of no more than 99%.

Polymorphic variant – A gene, mRNA, cDNA, polypeptide or peptide whose nucleotide or amino acid sequence varies from a reference sequence due to the presence of a polymorphism in the gene.

Polymorphism – The sequence variation observed in an individual at a polymorphic site. Polymorphisms include nucleotide substitutions, insertions, deletions and microsatellites and may, but need not, result in detectable differences in gene expression or protein function.

Polymorphism data – Information concerning one or more of the following for a specific gene: location of polymorphic sites; sequence variation at those sites; frequency of polymorphisms in one or more populations; the different genotypes and/or haplotypes determined for the gene; frequency of one or more of these genotypes and/or haplotypes in one or more populations; any known association(s) between a trait and a genotype or a haplotype for the gene.

Polymorphism Database – A collection of polymorphism data arranged in a systematic or methodical way and capable of being individually accessed by electronic or other means.

Polynucleotide – A nucleic acid molecule comprised of single-stranded RNA or DNA or comprised of complementary, double-stranded DNA.

Population Group – A group of individuals sharing a common ethnogeographic origin.

Reference Population – A group of subjects or individuals who are predicted to be representative of the genetic variation found in the general population. Typically, the reference population represents the genetic variation in the population at a certainty level of at least 85%, preferably at least 90%, more preferably at least 95% and even more preferably at least 99%.

Single Nucleotide Polymorphism (SNP) – Typically, the specific pair of nucleotides observed at a single polymorphic site. In rare cases, three or four nucleotides may be found.

Subject – A human individual whose genotypes or haplotypes or response to treatment or disease state are to be determined.

Treatment – A stimulus administered internally or externally to a subject.

Unphased – As applied to a sequence of nucleotide pairs for two or more polymorphic sites in a locus, unphased means the combination of nucleotides present at those polymorphic sites on a single copy of the locus is not known.

As discussed above, information on the identity of genotypes and haplotypes for the CTLA4 gene of any particular individual as well as the frequency of such genotypes and haplotypes in any particular population of individuals is expected to be useful for a variety of drug discovery and development applications. Thus, the invention also provides compositions and methods for detecting the novel CTLA4 polymorphisms and haplotypes identified herein.

The compositions comprise at least one CTLA4 genotyping oligonucleotide. In one embodiment, a CTLA4 genotyping oligonucleotide is a probe or primer capable of hybridizing to a target region that is located close to, or that contains, one of the novel polymorphic sites described herein. As used herein, the term "oligonucleotide" refers to a polynucleotide molecule having less than about 100 nucleotides. A preferred oligonucleotide of the invention is 10 to 35 nucleotides long. More preferably, the oligonucleotide is between 15 and 30, and most preferably, between 20 and 25 nucleotides in length. The exact length of the oligonucleotide will depend on many factors that are routinely considered and practiced by the skilled artisan. The oligonucleotide may be comprised of any phosphorylation state of ribonucleotides, deoxyribonucleotides, and acyclic nucleotide derivatives, and other functionally equivalent derivatives. Alternatively, oligonucleotides may have a phosphate-free backbone, which may be comprised of linkages such as carboxymethyl, acetamidate, carbamate, polyamide (peptide nucleic acid (PNA)) and the like (Varna, R. in Molecular Biology and Biotechnology, A Comprehensive Desk Reference, Ed. R. Meyers, VCH Publishers, Inc. (1995), pages 617-620). Oligonucleotides of the invention may be prepared by chemical synthesis using any suitable methodology known in the art, or may be derived from a biological sample, for example, by restriction digestion. The oligonucleotides may be labeled, according to any technique known in the art, including use of radiolabels, fluorescent labels, enzymatic labels, proteins, haptens, antibodies, sequence tags and the like.

Genotyping oligonucleotides of the invention must be capable of specifically hybridizing to a target region of a CTLA4 polynucleotide, i.e., a CTLA4 isogene. As used herein, specific hybridization means the oligonucleotide forms an anti-parallel double-stranded structure with the target region under certain hybridizing conditions, while failing to form such a structure when incubated with a non-target region or a non-CTLA4 polynucleotide under the same hybridizing conditions. Preferably, the oligonucleotide specifically hybridizes to the target region under conventional high stringency conditions. The skilled artisan can readily design and test oligonucleotide probes and primers suitable for detecting polymorphisms in the CTLA4 gene using the polymorphism information provided herein in conjunction with the known sequence information for the CTLA4 gene and routine techniques.

A nucleic acid molecule such as an oligonucleotide or polynucleotide is said to be a "perfect" or "complete" complement of another nucleic acid molecule if every nucleotide of one of the molecules is complementary to the nucleotide at the corresponding position of the other molecule. A nucleic acid molecule is "substantially complementary" to another molecule if it hybridizes to that molecule with sufficient stability to remain in a duplex form under conventional low-stringency

conditions. Conventional hybridization conditions are described, for example, by Sambrook J. et al., in *Molecular Cloning, A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Press, Cold Spring Harbor, NY (1989) and by Haymes, B.D. et al. in *Nucleic Acid Hybridization, A Practical Approach*, IRL Press, Washington, D.C. (1985). While perfectly complementary oligonucleotides are preferred for detecting polymorphisms, departures from complete complementarity are contemplated where such departures do not prevent the molecule from specifically hybridizing to the target region. For example, an oligonucleotide primer may have a non-complementary fragment at its 5' end, with the remainder of the primer being complementary to the target region. Alternatively, non-complementary nucleotides may be interspersed into the oligonucleotide probe or primer as long as the resulting probe or primer is still capable of specifically hybridizing to the target region.

Preferred genotyping oligonucleotides of the invention are allele-specific oligonucleotides. As used herein, the term allele-specific oligonucleotide (ASO) means an oligonucleotide that is able, under sufficiently stringent conditions, to hybridize specifically to one allele of a gene, or other locus, at a target region containing a polymorphic site while not hybridizing to the corresponding region in another allele(s). As understood by the skilled artisan, allele-specificity will depend upon a variety of readily optimized stringency conditions, including salt and formamide concentrations, as well as temperatures for both the hybridization and washing steps. Examples of hybridization and washing conditions typically used for ASO probes are found in Kogan et al., "Genetic Prediction of Hemophilia A" in *PCR Protocols, A Guide to Methods and Applications*, Academic Press, 1990 and Ruaño et al., 87 *Proc. Natl. Acad. Sci. USA* 6296-6300, 1990. Typically, an ASO will be perfectly complementary to one allele while containing a single mismatch for another allele.

Allele-specific oligonucleotides of the invention include ASO probes and ASO primers. ASO probes which usually provide good discrimination between different alleles are those in which a central position of the oligonucleotide probe aligns with the polymorphic site in the target region (e.g., approximately the 7th or 8th position in a 15mer, the 8th or 9th position in a 16mer, and the 10th or 11th position in a 20mer). An ASO primer of the invention has a 3' terminal nucleotide, or preferably a 3' penultimate nucleotide, that is complementary to only one nucleotide of a particular SNP, thereby acting as a primer for polymerase-mediated extension only if the allele containing that nucleotide is present. ASO probes and primers hybridizing to either the coding or noncoding strand are contemplated by the invention.

ASO probes and primers listed below use the appropriate nucleotide symbol (R= G or A, Y= T or C, M= A or C, K= G or T, S= G or C, and W= A or T; WIPO standard ST.25) at the position of the polymorphic site to represent the two alternative allelic variants observed at that polymorphic site.

A preferred ASO probe for detecting CTLA4 gene polymorphisms comprises a nucleotide sequence, listed 5' to 3', selected from the group consisting of:

AGATCCTYAAAGTGA	(SEQ ID NO:4) and its complement,
CAGTCAARGGCAGTG	(SEQ ID NO:5) and its complement,
CACTGAGYTGACACC	(SEQ ID NO:6) and its complement,
CTAGAACYGTAGGCA	(SEQ ID NO:7) and its complement,

TTTTAATRGCTGAAT (SEQ ID NO:9) and its complement, and
GCTGTGARCATTTCAT (SEQ ID NO:9) and its complement.

A preferred ASO primer for detecting CTLA4 gene polymorphisms comprises a nucleotide sequence, listed 5' to 3', selected from the group consisting of:

TTATCCAGATCCTYA (SEQ ID NO:10); TCATGTTCACTTTTRA (SEQ ID NO:11);
TTTCAGCAGTCAARG (SEQ ID NO:12); ATAAATCACTGCCYT (SEQ ID NO:13);
CCATTTCAGTGAGYT (SEQ ID NO:14); GCAACAGGTGTCARC (SEQ ID NO:15);
AACGCACTAGAACYG (SEQ ID NO:16); TGCCAATGCCTACRG (SEQ ID NO:17);
AATAAATTTTAATRG (SEQ ID NO:18); TTCTTGATTTCAGCYA (SEQ ID NO:19);
CTGTATGCTGTGARC (SEQ ID NO:20); and TTAAAAATGAATGYT (SEQ ID NO:21).

Other genotyping oligonucleotides of the invention hybridize to a target region located one to several nucleotides downstream of one of the novel polymorphic sites identified herein. Such oligonucleotides are useful in polymerase-mediated primer extension methods for detecting one of the novel polymorphisms described herein and therefore such genotyping oligonucleotides are referred to herein as "primer-extension oligonucleotides". In a preferred embodiment, the 3'-terminus of a primer-extension oligonucleotide is a deoxynucleotide complementary to the nucleotide located immediately adjacent to the polymorphic site.

A particularly preferred oligonucleotide primer for detecting CTLA4 gene polymorphisms by primer extension terminates in a nucleotide sequence, listed 5' to 3', selected from the group consisting of:

TCCAGATCCT (SEQ ID NO:22); TGTTCACTTT (SEQ ID NO:23);
CAGCAGTCAA (SEQ ID NO:24); AATCACTGCC (SEQ ID NO:25);
TTTCACTGAG (SEQ ID NO:26); ACAGGTGTCA (SEQ ID NO:27);
GCACTAGAAC (SEQ ID NO:28); CAATGCCTAC (SEQ ID NO:29);
AAATTTTAAT (SEQ ID NO:30); TTGATTCAGC (SEQ ID NO:31);
TATGCTGTGA (SEQ ID NO:32); and AAAATGAATG (SEQ ID NO:33).

In some embodiments, a composition contains two or more differently labeled genotyping oligonucleotides for simultaneously probing the identity of nucleotides at two or more polymorphic sites. It is also contemplated that primer compositions may contain two or more sets of allele-specific primer pairs to allow simultaneous targeting and amplification of two or more regions containing a polymorphic site.

CTLA4 genotyping oligonucleotides of the invention may also be immobilized on or synthesized on a solid surface such as a microchip, bead, or glass slide (see, e.g., WO 98/20020 and WO 98/20019). Such immobilized genotyping oligonucleotides may be used in a variety of polymorphism detection assays, including but not limited to probe hybridization and polymerase extension assays. Immobilized CTLA4 genotyping oligonucleotides of the invention may comprise an ordered array of oligonucleotides designed to rapidly screen a DNA sample for polymorphisms in multiple genes at the same time.

individual is heterozygous for that site. The polymorphism may be identified directly, known as positive-type identification, or by inference, referred to as negative-type identification. For example, where a SNP is known to be guanine and cytosine in a reference population, a site may be positively determined to be either guanine or cytosine for an individual homozygous at that site, or both guanine and cytosine, if the individual is heterozygous at that site. Alternatively, the site may be negatively determined to be not guanine (and thus cytosine/cytosine) or not cytosine (and thus guanine/guanine).

The target region(s) may be amplified using any oligonucleotide-directed amplification method, including but not limited to polymerase chain reaction (PCR) (U.S. Patent No. 4,965,188), ligase chain reaction (LCR) (Barany et al., *Proc. Natl. Acad. Sci. USA* 88:189-193, 1991; WO90/01069), and oligonucleotide ligation assay (OLA) (Landegren et al., *Science* 241:1077-1080, 1988).

Other known nucleic acid amplification procedures may be used to amplify the target region including transcription-based amplification systems (U.S. Patent No. 5,130,238; EP 329,822; U.S. Patent No. 5,169,766, WO89/06700) and isothermal methods (Walker et al., *Proc. Natl. Acad. Sci. USA* 89:392-396, 1992).

A polymorphism in the target region may also be assayed before or after amplification using one of several hybridization-based methods known in the art. Typically, allele-specific oligonucleotides are utilized in performing such methods. The allele-specific oligonucleotides may be used as differently labeled probe pairs, with one member of the pair showing a perfect match to one variant of a target sequence and the other member showing a perfect match to a different variant. In some embodiments, more than one polymorphic site may be detected at once using a set of allele-specific oligonucleotides or oligonucleotide pairs. Preferably, the members of the set have melting temperatures within 5°C, and more preferably within 2°C, of each other when hybridizing to each of the polymorphic sites being detected.

Hybridization of an allele-specific oligonucleotide to a target polymucleotide may be performed with both entities in solution, or such hybridization may be performed when either the oligonucleotide or the target polymucleotide is covalently or noncovalently affixed to a solid support. Attachment may be mediated, for example, by antibody-antigen interactions, poly-L-Lys, streptavidin or avidin-biotin, salt bridges, hydrophobic interactions, chemical linkages, UV cross-linking baking, etc. Allele-specific oligonucleotides may be synthesized directly on the solid support or attached to the solid support subsequent to synthesis. Solid-supports suitable for use in detection methods of the invention include substrates made of silicon, glass, plastic, paper and the like, which may be formed, for example, into wells (as in 96-well plates), slides, sheets, membranes, fibers, chips, dishes, and beads. The solid support may be treated, coated or derivatized to facilitate the immobilization of the allele-specific oligonucleotide or target nucleic acid.

The genotype or haplotype for the CTLA4 gene of an individual may also be determined by hybridization of a nucleic acid sample containing one or both copies of the gene, or fragment(s) thereof, to nucleic acid arrays and subarrays such as described in WO 95/11995. The arrays would

contain a battery of allele-specific oligonucleotides representing each of the polymorphic sites to be included in the genotype or haplotype.

The identity of polymorphisms may also be determined using a mismatch detection technique, including but not limited to the RNase protection method using riboprobes (Winter et al., *Proc. Natl. Acad. Sci. USA* 82:7575, 1985; Meyers et al., *Science* 230:1242, 1985) and proteins which recognize nucleotide mismatches, such as the *E. coli* mutS protein (Modrich, *P. Ann. Rev. Genet.* 25:229-253, 1991). Alternatively, variant alleles can be identified by single strand conformation polymorphism (SSCP) analysis (Orita et al., *Genomics* 5:874-879, 1989; Humphries et al., in *Molecular Diagnosis of Genetic Diseases*, R. Elles, ed., pp. 321-340, 1996) or denaturing gradient gel electrophoresis (DGGE) (Wartell et al., *Nucl. Acids Res.* 18:2699-2706, 1990; Sheffield et al., *Proc. Natl. Acad. Sci. USA* 86:232-236, 1989).

A polymerase-mediated primer extension method may also be used to identify the polymorphism(s). Several such methods have been described in the patent and scientific literature and include the "Genetic Bit Analysis" method (WO92/15712) and the ligase/polymerase mediated genetic bit analysis (U.S. Patent 5,679,524. Related methods are disclosed in WO91/02087, WO90/09455, WO95/17676, U.S. Patent Nos. 5,302,509, and 5,945,283. Extended primers containing a polymorphism may be detected by mass spectrometry as described in U.S. Patent No. 5,605,798. Another primer extension method is allele-specific PCR (Ruano et al., *Nucl. Acids Res.* 17:8392, 1989; Ruano et al., *Nucl. Acids Res.* 19, 6877-6882, 1991; WO 93/22456; Turki et al., *J. Clin. Invest.* 95:1635-1641, 1995). In addition, multiple polymorphic sites may be investigated by simultaneously amplifying multiple regions of the nucleic acid using sets of allele-specific primers as described in Wallace et al. (WO89/10414).

In addition, the identity of the allele(s) present at any of the novel polymorphic sites described herein may be indirectly determined by genotyping another polymorphic site that is in linkage disequilibrium with the polymorphic site that is of interest. Polymorphic sites in linkage disequilibrium with the presently disclosed polymorphic sites may be located in regions of the gene or in other genomic regions not examined herein. Genotyping of a polymorphic site in linkage disequilibrium with the novel polymorphic sites described herein may be performed by, but is not limited to, any of the above-mentioned methods for detecting the identity of the allele at a polymorphic site.

In another aspect of the invention, an individual's CTLA4 haplotype pair is predicted from its CTLA4 genotype using information on haplotype pairs known to exist in a reference population. In its broadest embodiment, the haplotyping prediction method comprises identifying a CTLA4 genotype for the individual at two or more CTLA4 polymorphic sites described herein, enumerating all possible haplotype pairs which are consistent with the genotype, accessing data containing CTLA4 haplotype pairs identified in a reference population, and assigning a haplotype pair to the individual that is consistent with the data. In one embodiment, the reference haplotype pairs include the CTLA4 haplotype pairs shown in Table 3.

Generally, the reference population should be composed of randomly-selected individuals representing the major ethnogeographic groups of the world. A preferred reference population for use in the methods of the present invention comprises an approximately equal number of individuals from Caucasian, African-descent, Asian and Hispanic-Latino population groups with the minimum number of each group being chosen based on how rare a haplotype one wants to be guaranteed to see. For example, if one wants to have a $q\%$ chance of not missing a haplotype that exists in the population at a $p\%$ frequency of occurring in the reference population, the number of individuals (n) who must be sampled is given by $2n = \log(1-q)/\log(1-p)$ where p and q are expressed as fractions. A preferred reference population allows the detection of any haplotype whose frequency is at least 10% with about 99% certainty and comprises about 20 unrelated individuals from each of the four population groups named above. A particularly preferred reference population includes a 3-generation family representing one or more of the four population groups to serve as controls for checking quality of haplotyping procedures.

In a preferred embodiment, the haplotype frequency data for each ethnogeographic group is examined to determine whether it is consistent with Hardy-Weinberg equilibrium. Hardy-Weinberg equilibrium (D.L. Hartl et al., *Principles of Population Genomics*, Sinauer Associates (Sunderland, MA), 3rd Ed., 1997) postulates that the frequency of finding the haplotype pair H_1 / H_2 is equal to $p_{H-W}(H_1 / H_2) = 2p(H_1)p(H_2)$ if $H_1 \neq H_2$ and $p_{H-W}(H_1 / H_2) = p(H_1)p(H_2)$ if $H_1 = H_2$. A statistically significant difference between the observed and expected haplotype frequencies could be due to one or more factors including significant inbreeding in the population group, strong selective pressure on the gene, sampling bias, and/or errors in the genotyping process. If large deviations from Hardy-Weinberg equilibrium are observed in an ethnogeographic group, the number of individuals in that group can be increased to see if the deviation is due to a sampling bias. If a larger sample size does not reduce the difference between observed and expected haplotype pair frequencies, then one may wish to consider haplotyping the individual using a direct haplotyping method such as, for example, CLASPER System™ technology (U.S. Patent No. 5,866,404), single molecule dilution, or allele-specific long-range PCR (Michalotos-Beloin et al., *Nucleic Acids Res.* 24:4841-4843, 1996).

In one embodiment of this method for predicting a CTLA4 haplotype pair for an individual, the assigning step involves performing the following analysis. First, each of the possible haplotype pairs is compared to the haplotype pairs in the reference population. Generally, only one of the haplotype pairs in the reference population matches a possible haplotype pair and that pair is assigned to the individual. Occasionally, only one haplotype represented in the reference haplotype pairs is consistent with a possible haplotype pair for an individual, and in such cases the individual is assigned a haplotype pair containing this known haplotype and a new haplotype derived by subtracting the known haplotype from the possible haplotype pair. In rare cases, either no haplotypes in the reference population are consistent with the possible haplotype pairs, or alternatively, multiple reference haplotype pairs are consistent with the possible haplotype pairs. In such cases, the individual is

diseases and other disorders. As used herein the term "clinical response" means any or all of the following: a quantitative measure of the response, no response, and adverse response (i.e., side effects).

In order to deduce a correlation between clinical response to a treatment and a CTLA4 genotype, haplotype, or haplotype pair, it is necessary to obtain data on the clinical responses exhibited by a population of individuals who received the treatment, hereinafter the "clinical population". This clinical data may be obtained by analyzing the results of a clinical trial that has already been run and/or the clinical data may be obtained by designing and carrying out one or more new clinical trials. As used herein, the term "clinical trial" means any research study designed to collect clinical data on responses to a particular treatment, and includes but is not limited to phase I, phase II and phase III clinical trials. Standard methods are used to define the patient population and to enroll subjects.

It is preferred that the individuals included in the clinical population have been graded for the existence of the medical condition of interest. This is important in cases where the symptom(s) being presented by the patients can be caused by more than one underlying condition, and where treatment of the underlying conditions are not the same. An example of this would be where patients experience breathing difficulties that are due to either asthma or respiratory infections. If both sets were treated with an asthma medication, there would be a spurious group of apparent non-responders that did not actually have asthma. These people would affect the ability to detect any correlation between haplotype and treatment outcome. This grading of potential patients could employ a standard physical exam or one or more lab tests. Alternatively, grading of patients could use haplotyping for situations where there is a strong correlation between haplotype pair and disease susceptibility or severity.

The therapeutic treatment of interest is administered to each individual in the trial population and each individual's response to the treatment is measured using one or more predetermined criteria. It is contemplated that in many cases, the trial population will exhibit a range of responses and that the investigator will choose the number of responder groups (e.g., low, medium, high) made up by the various responses. In addition, the CTLA4 gene for each individual in the trial population is genotyped and/or haplotyped, which may be done before or after administering the treatment.

After both the clinical and polymorphism data have been obtained, correlations between individual response and CTLA4 genotype or haplotype content are created. Correlations may be produced in several ways. In one method, individuals are grouped by their CTLA4 genotype or haplotype (or haplotype pair) (also referred to as a polymorphism group), and then the averages and standard deviations of clinical responses exhibited by the members of each polymorphism group are calculated.

These results are then analyzed to determine if any observed variation in clinical response between polymorphism groups is statistically significant. Statistical analysis methods which may be used are described in L.D. Fisher and G. vanBelle, "Biostatistics: A Methodology for the Health Sciences", Wiley-Interscience (New York) 1993. This analysis may also include a regression calculation of which polymorphic sites in the CTLA4 gene give the most significant contribution to the differences in phenotype. One regression model useful in the invention is described in PCT

Application Serial No. PCT/US00/17540, entitled "Methods for Obtaining and Using Haplotype Data".

A second method for finding correlations between CTLA4 haplotype content and clinical responses uses predictive models based on error-minimizing optimization algorithms. One of many possible optimization algorithms is a genetic algorithm (R. Judson, "Genetic Algorithms and Their Uses in Chemistry" in Reviews in Computational Chemistry, Vol. 10, pp. 1-73, K. B. Lipkowitz and D. B. Boyd, eds. (VCH Publishers, New York, 1997). Simulated annealing (Press et al., "Numerical Recipes in C: The Art of Scientific Computing", Cambridge University Press (Cambridge) 1992, Ch. 10), neural networks (E. Rich and K. Knight, "Artificial Intelligence", 2nd Edition (McGraw-Hill, New York, 1991, Ch. 18), standard gradient descent methods (Press et al., *supra*, Ch. 10), or other global or local optimization approaches (see discussion in Judson, *supra*) could also be used. Preferably, the correlation is found using a genetic algorithm approach as described in PCT Application Serial No. PCT/US00/17540.

Correlations may also be analyzed using analysis of variation (ANOVA) techniques to determine how much of the variation in the clinical data is explained by different subsets of the polymorphic sites in the CTLA4 gene. As described in PCT Application Serial No. PCT/US00/17540, ANOVA is used to test hypotheses about whether a response variable is caused by or correlated with one or more traits or variables that can be measured (Fisher and vanBelle, *supra*, Ch. 10).

From the analyses described above, a mathematical model may be readily constructed by the skilled artisan that predicts clinical response as a function of CTLA4 genotype or haplotype content. Preferably, the model is validated in one or more follow-up clinical trials designed to test the model.

The identification of an association between a clinical response and a genotype or haplotype (or haplotype pair) for the CTLA4 gene may be the basis for designing a diagnostic method to determine those individuals who will or will not respond to the treatment, or alternatively, will respond at a lower level and thus may require more treatment, i.e., a greater dose of a drug. The diagnostic method may take one of several forms: for example, a direct DNA test (i.e., genotyping or haplotyping one or more of the polymorphic sites in the CTLA4 gene), a serological test, or a physical exam measurement. The only requirement is that there be a good correlation between the diagnostic test results and the underlying CTLA4 genotype or haplotype that is in turn correlated with the clinical response. In a preferred embodiment, this diagnostic method uses the predictive haplotyping method described above.

In another embodiment, the invention provides an isolated polynucleotide comprising a polymorphic variant of the CTLA4 gene or a fragment of the gene which contains at least one of the novel polymorphic sites described herein. The nucleotide sequence of a variant CTLA4 gene is identical to the reference genomic sequence for those portions of the gene examined, as described in the Examples below, except that it comprises a different nucleotide at one or more of the novel polymorphic sites PS1, PS3, PS4, PS5, PS6 and PS7, and may also comprise an additional polymorphism of guanine at PS2. Similarly, the nucleotide sequence of a variant fragment of the

CTLA4 gene is identical to the corresponding portion of the reference sequence except for having a different nucleotide at one or more of the novel polymorphic sites described herein. Thus, the invention specifically does not include polynucleotides comprising a nucleotide sequence identical to the reference sequence of the CTLA4 gene, which is defined by haplotype 1, (or other reported CTLA4 sequences) or to portions of the reference sequence (or other reported CTLA4 sequences), except for genotyping oligonucleotides as described above.

The location of a polymorphism in a variant gene or fragment is identified by aligning its sequence against SEQ ID NO:1. The polymorphism is selected from the group consisting of thymine at PS1, guanine at PS3, cytosine at PS4, thymine at PS5, guanine at PS6 and guanine at PS7. In a preferred embodiment, the polymorphic variant comprises a naturally-occurring isogene of the CTLA4 gene which is defined by any one of haplotypes 2-8 shown in Table 4 below.

Polymorphic variants of the invention may be prepared by isolating a clone containing the CTLA4 gene from a human genomic library. The clone may be sequenced to determine the identity of the nucleotides at the novel polymorphic sites described herein. Any particular variant claimed herein could be prepared from this clone by performing *in vitro* mutagenesis using procedures well-known in the art.

CTLA4 isogenes may be isolated using any method that allows separation of the two "copies" of the CTLA4 gene present in an individual, which, as readily understood by the skilled artisan, may be the same allele or different alleles. Separation methods include targeted *in vivo* cloning (TIVC) in yeast as described in WO 98/01573, U.S. Patent No. 5,866,404, and U.S. Patent No. 5,972,614. Another method, which is described in U.S. Patent No. 5,972,614, uses an allele specific oligonucleotide in combination with primer extension and exonuclease degradation to generate hemizygous DNA targets. Yet other methods are single molecule dilution (SMD) as described in Ruaño et al., *Proc. Natl. Acad. Sci.* 87:6296-6300, 1990; and allele specific PCR (Ruaño et al., 1989, *supra*; Ruaño et al., 1991, *supra*; Michalatos-Beloin et al., *supra*).

The invention also provides CTLA4 genome anthologies, which are collections of CTLA4 isogenes found in a given population. The population may be any group of at least two individuals, including but not limited to a reference population, a population group, a family population, a clinical population, and a same sex population. A CTLA4 genome anthology may comprise individual CTLA4 isogenes stored in separate containers such as microtest tubes, separate wells of a microtitre plate and the like. Alternatively, two or more groups of the CTLA4 isogenes in the anthology may be stored in separate containers. Individual isogenes or groups of isogenes in a genome anthology may be stored in any convenient and stable form, including but not limited to in buffered solutions, as DNA precipitates, freeze-dried preparations and the like. A preferred CTLA4 genome anthology of the invention comprises a set of isogenes defined by the haplotypes shown in Table 4 below.

An isolated polynucleotide containing a polymorphic variant nucleotide sequence of the invention may be operably linked to one or more expression regulatory elements in a recombinant expression vector capable of being propagated and expressing the encoded CTLA4 protein in a

prokaryotic or a eukaryotic host cell. Examples of expression regulatory elements which may be used include, but are not limited to, the lac system, operator and promoter regions of phage lambda, yeast promoters, and promoters derived from vaccinia virus, adenovirus, retroviruses, or SV40. Other regulatory elements include, but are not limited to, appropriate leader sequences, termination codons, polyadenylation signals, and other sequences required for the appropriate transcription and subsequent translation of the nucleic acid sequence in a given host cell. Of course, the correct combinations of expression regulatory elements will depend on the host system used. In addition, it is understood that the expression vector contains any additional elements necessary for its transfer to and subsequent replication in the host cell. Examples of such elements include, but are not limited to, origins of replication and selectable markers. Such expression vectors are commercially available or are readily constructed using methods known to those in the art (e.g., F. Ausubel et al., 1987, in "Current Protocols in Molecular Biology", John Wiley and Sons, New York, New York). Host cells which may be used to express the variant CTLA4 sequences of the invention include, but are not limited to, eukaryotic and mammalian cells, such as animal, plant, insect and yeast cells, and prokaryotic cells, such as *E. coli*, or algal cells as known in the art. The recombinant expression vector may be introduced into the host cell using any method known to those in the art including, but not limited to, microinjection, electroporation, particle bombardment, transduction, and transfection using DEAE-dextran, lipofection, or calcium phosphate (see e.g., Sambrook et al. (1989) in "Molecular Cloning. A Laboratory Manual", Cold Spring Harbor Press, Plainview, New York). In a preferred aspect, eukaryotic expression vectors that function in eukaryotic cells, and preferably mammalian cells, are used. Non-limiting examples of such vectors include vaccinia virus vectors, adenovirus vectors, herpes virus vectors, and baculovirus transfer vectors. Preferred eukaryotic cell lines include COS cells, CHO cells, HeLa cells, NIH/3T3 cells, and embryonic stem cells (Thomson, J. A. et al., 1998 *Science* 282:1145-1147). Particularly preferred host cells are mammalian cells.

As used herein, a polymorphic variant of a CTLA4 gene fragment comprises at least one novel polymorphism identified herein and has a length of at least 10 nucleotides and may range up to the full length of the gene. Preferably, such fragments are between 100 and 3000 nucleotides in length, and more preferably between 200 and 2000 nucleotides in length, and most preferably between 500 and 1000 nucleotides in length.

In describing the CTLA4 polymorphic sites identified herein, reference is made to the sense strand of the gene for convenience. However, as recognized by the skilled artisan, nucleic acid molecules containing the CTLA4 gene may be complementary double stranded molecules and thus reference to a particular site on the sense strand refers as well to the corresponding site on the complementary antisense strand. Thus, reference may be made to the same polymorphic site on either strand and an oligonucleotide may be designed to hybridize specifically to either strand at a target region containing the polymorphic site. Thus, the invention also includes single-stranded polynucleotides which are complementary to the sense strand of the CTLA4 genomic variants described herein.

Polynucleotides comprising a polymorphic gene variant or fragment may be useful for therapeutic purposes. For example, where a patient could benefit from expression, or increased expression, of a particular CTLA4 protein isoform, an expression vector encoding the isoform may be administered to the patient. The patient may be one who lacks the CTLA4 isogene encoding that isoform or may already have at least one copy of that isogene.

In other situations, it may be desirable to decrease or block expression of a particular CTLA4 isogene. Expression of a CTLA4 isogene may be turned off by transforming a targeted organ, tissue or cell population with an expression vector that expresses high levels of untranslatable mRNA for the isogene. Alternatively, oligonucleotides directed against the regulatory regions (e.g., promoter, introns, enhancers, 3' untranslated region) of the isogene may block transcription. Oligonucleotides targeting the transcription initiation site, e.g., between positions -10 and +10 from the start site are preferred. Similarly, inhibition of transcription can be achieved using oligonucleotides that base-pair with region(s) of the isogene DNA to form triplex DNA (see e.g., Gee et al. in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, N.Y., 1994). Antisense oligonucleotides may also be designed to block translation of CTLA4 mRNA transcribed from a particular isogene. It is also contemplated that ribozymes may be designed that can catalyze the specific cleavage of CTLA4 mRNA transcribed from a particular isogene.

The oligonucleotides may be delivered to a target cell or tissue by expression from a vector introduced into the cell or tissue *in vivo* or *ex vivo*. Alternatively, the oligonucleotides may be formulated as a pharmaceutical composition for administration to the patient. Oligoribonucleotides and/or oligodeoxynucleotides intended for use as antisense oligonucleotides may be modified to increase stability and half-life. Possible modifications include, but are not limited to phosphorothioate or 2' O-methyl linkages, and the inclusion of nontraditional bases such as inosine and queosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytosine, guanine, thymine, and uracil which are not as easily recognized by endogenous nucleases.

Effect(s) of the polymorphisms identified herein on expression of CTLA4 may be investigated by preparing recombinant cells and/or nonhuman recombinant organisms, preferably recombinant animals, containing a polymorphic variant of the CTLA4 gene. As used herein, "expression" includes but is not limited to one or more of the following: transcription of the gene into precursor mRNA; splicing and other processing of the precursor mRNA to produce mature mRNA; mRNA stability; translation of the mature mRNA into CTLA4 protein (including codon usage and tRNA availability); and glycosylation and/or other modifications of the translation product, if required for proper expression and function.

To prepare a recombinant cell of the invention, the desired CTLA4 isogene may be introduced into the cell in a vector such that the isogene remains extrachromosomal. In such a situation, the gene will be expressed by the cell from the extrachromosomal location. In a preferred embodiment, the CTLA4 isogene is introduced into a cell in such a way that it recombines with the endogenous CTLA4 gene present in the cell. Such recombination requires the occurrence of a double recombination event,

thereby resulting in the desired CTLA4 gene polymorphism. Vectors for the introduction of genes both for recombination and for extrachromosomal maintenance are known in the art, and any suitable vector or vector construct may be used in the invention. Methods such as electroporation, particle bombardment, calcium phosphate co-precipitation and viral transduction for introducing DNA into cells are known in the art; therefore, the choice of method may lie with the competence and preference of the skilled practitioner. Examples of cells into which the CTLA4 isogene may be introduced include, but are not limited to, continuous culture cells, such as COS, NIH/3T3, and primary or culture cells of the relevant tissue type, i.e., they express the CTLA4 isogene. Such recombinant cells can be used to compare the biological activities of the different protein variants.

Recombinant nonhuman organisms, i.e., transgenic animals, expressing a variant CTLA4 gene are prepared using standard procedures known in the art. Preferably, a construct comprising the variant gene is introduced into a nonhuman animal or an ancestor of the animal at an embryonic stage, i.e., the one-cell stage, or generally not later than about the eight-cell stage. Transgenic animals carrying the constructs of the invention can be made by several methods known to those having skill in the art. One method involves transfecting into the embryo a retrovirus constructed to contain one or more insulator elements, a gene or genes of interest, and other components known to those skilled in the art to provide a complete shuttle vector harboring the insulated gene(s) as a transgene, see e.g., U.S. Patent No. 5,610,053. Another method involves directly injecting a transgene into the embryo. A third method involves the use of embryonic stem cells. Examples of animals into which the CTLA4 isogenes may be introduced include, but are not limited to, mice, rats, other rodents, and nonhuman primates (see "The Introduction of Foreign Genes into Mice" and the cited references therein, In: Recombinant DNA, Eds. J.D. Watson, M. Gilman, J. Witkowski, and M. Zoller, W.H. Freeman and Company, New York, pages 254-272). Transgenic animals stably expressing a human CTLA4 isogen and producing human CTLA4 protein can be used as biological models for studying diseases related to abnormal CTLA4 expression and/or activity, and for screening and assaying various candidate drugs, compounds, and treatment regimens to reduce the symptoms or effects of these diseases.

An additional embodiment of the invention relates to pharmaceutical compositions for treating disorders affected by expression or function of a novel CTLA4 isogene described herein. The pharmaceutical composition may comprise any of the following active ingredients: a polynucleotide comprising one of these novel CTLA4 isogenes; an antisense oligonucleotide directed against one of the novel CTLA4 isogenes, a polynucleotide encoding such an antisense oligonucleotide, or another compound which inhibits expression of a novel CTLA4 isogene described herein. Preferably, the composition contains the active ingredient in a therapeutically effective amount. By therapeutically effective amount is meant that one or more of the symptoms relating to disorders affected by expression or function of a novel CTLA4 isogene is reduced and/or eliminated. The composition also comprises a pharmaceutically acceptable carrier, examples of which include, but are not limited to, saline, buffered saline, dextrose, and water. Those skilled in the art may employ a formulation most suitable for the active ingredient, whether it is a polynucleotide, oligonucleotide, protein, peptide or

isolation, PCR and sequencing procedures. Such methods are well-known to those skilled in the art and are described in numerous publications, for example, Sambrook, Fritsch, and Maniatis, "Molecular Cloning: A Laboratory Manual", 2nd Edition, Cold Spring Harbor Laboratory Press, USA, (1989).

EXAMPLE 1

This example illustrates examination of various regions of the CTLA4 gene for polymorphic sites.

Amplification of Target Regions

The following target regions were amplified using either the PCR primers represented below or 'tailed' PCR primers, each of which includes a universal sequence forming a noncomplementary 'tail' attached to the 5' end of each unique sequence in the PCR primer pairs. The universal 'tail' sequence for the forward PCR primers comprises the sequence 5'-TGTAACGACGGCCAGT-3' (SEQ ID NO:34) and the universal 'tail' sequence for the reverse PCR primers comprises the sequence 5'-AGGAAACAGCTATGACCAT-3' (SEQ ID NO:35). The nucleotide positions of the first and last nucleotide of the forward and reverse primers for each region amplified are presented below and correspond to positions in Figure 1.

PCR Primer Pairs

Fragment No.	Forward Primer	Reverse Primer	PCR Product
Fragment 1	37286-37308	complement of 37909-37889	624 nt
Fragment 2	37608-37630	complement of 38295-38273	688 nt
Fragment 3	37690-37711	complement of 38108-38088	419 nt
Fragment 4	40310-40333	complement of 40820-40798	511 nt
Fragment 5	40466-40488	complement of 41010-40988	545 nt
Fragment 6	41123-41147	complement of 41644-41621	522 nt
Fragment 7	42294-42315	complement of 42918-42896	625 nt

These primer pairs were used in PCR reactions containing genomic DNA isolated from immortalized cell lines for each member of the Index Repository. The PCR reactions were carried out under the following conditions:

Reaction volume	= 10 µl
10 x Advantage 2 Polymerase reaction buffer (Clontech)	= 1 µl
100 ng of human genomic DNA	= 1 µl
10 mM dNTP	= 0.4 µl
Advantage 2 Polymerase enzyme mix (Clontech)	= 0.2 µl
Forward Primer (10 µM)	= 0.4 µl
Reverse Primer (10 µM)	= 0.4 µl
Water	= 6.6 µl

Amplification profile:

97°C - 2 min. 1 cycle

97°C - 15 sec.

}

70°C - 45 sec. 10 cycles
72°C - 45 sec.

97°C - 15 sec.
64°C - 45 sec. } 35 cycles
72°C - 45 sec.

Sequencing of PCR Products

The PCR products were purified using a Whatman/Polyfritronics 100 µl 384 well unfilter plate essentially according to the manufacturers protocol. The purified DNA was eluted in 50 µl of distilled water. Sequencing reactions were set up using Applied Biosystems Big Dye Terminator chemistry essentially according to the manufacturers protocol. The purified PCR products were sequenced in both directions using either the primer sets represented below with the positions of their first and last nucleotide corresponding to positions in Figure 1, or the appropriate universal 'tail' sequence as a primer. Reaction products were purified by isopropanol precipitation, and run on an Applied Biosystems 3700 DNA Analyzer.

Sequencing Primer Pairs

Fragment No.	Forward Primer	Reverse Primer
Fragment 1	37358-37377	complement of 37882-37863
Fragment 2	37637-37656	complement of 38175-38155
Fragment 3	Tailed Seq	
Fragment 4	40349-40368	complement of 40789-40769
Fragment 5	Tailed Seq.	
Fragment 6	41162-41182	complement of 41565-41546
Fragment 7	42340-42361	complement of 42868-42850

Analysis of Sequences for Polymorphic Sites

Sequence information for a minimum of 80 humans was analyzed for the presence of polymorphisms using the Polyphred program (Nickerson et al., *Nucleic Acids Res.* 14:2745-2751, 1997). The presence of a polymorphism was confirmed on both strands. The polymorphisms and their locations in the CTLA4 gene are listed in Table 2 below.

Table 2. Polymorphic Sites Identified in the CTLA4 Gene

Polymorphic Site Number	PolyId ^a	Nucleotide Position	Reference Allele	Variant Allele	CDS Variant	AA Variant
PS1	743786	37535	C	T		
PS2 ^a	743788	37902	A	G	49	T17A
PS3	743794	38038	A	G		
PS4	743804	40867	T	C		
PS5	743808	41547	C	T		
PS6	743812	42460	A	G		
PS7	743814	42508	A	G		

^aPolyId is a unique identifier assigned to each PS by Genassance Pharmaceuticals, Inc.

^bPreviously identified in literature

EXAMPLE 2

This example illustrates analysis of the CTLA4 polymorphisms identified in the Index

Repository for human genotypes and haplotypes.

The different genotypes containing these polymorphisms that were observed in the reference population are shown in Table 3 below, with the haplotype pair indicating the combination of haplotypes determined for the individual using the haplotype derivation protocol described below. In Table 3, homozygous positions are indicated by one nucleotide and heterozygous positions are indicated by two nucleotides. Missing nucleotides in any given genotype in Table 3 were inferred based on linkage disequilibrium and/or Mendelian inheritance.

Table 3. Genotypes and Haplotype Pairs Observed for CTLA4 Gene

Genotype Number	Polymorphic Sites							HAP Pair	
	PS1	PS2	PS3	PS4	PS5	PS6	PS7		
1	C	A	G	T	C	A	A	2	2
2	C	G	G	T	C	A	A	7	7
3	C	A	G/A	T	C	A	A	2	1
4	C	A	G	T	C	A	A/G	2	3
5	C	A	G	T	C/T	A	A	2	5
6	C	A/G	G	T/C	C	A	A	2	6
7	C/T	A	G	T	C	A	A	2	8
8	C	A	G	T	C	A/G	A	2	4
9	C/T	G/A	G	T	C	A	A	7	8
10	C	A/G	G	T	C	A	A	2	7

The haplotype pairs shown in Table 3 were estimated from the unphased genotypes using a computer-implemented extension of Clark's algorithm (Clark, A.G. 1990 *Mol Bio Evol* 7, 111-122) for assigning haplotypes to unrelated individuals in a population sample, as described in U.S. Provisional Application Serial No. 60/198,340 entitled "A Method and System for Determining Haplotypes from a Collection of Polymorphisms" and the corresponding International Application filed April 18, 2001. In this method, haplotypes are assigned directly from individuals who are homozygous at all sites or heterozygous at no more than one of the variable sites. This list of haplotypes is augmented with haplotypes obtained from two families (one three-generation Caucasian family and one two-generation African-American family) and then used to deconvolute the unphased genotypes in the remaining (multiply heterozygous) individuals.

By following this protocol, it was determined that the Index Repository examined herein and, by extension, the general population contains the 8 human CTLA4 haplotypes shown in Table 4 below.

Table 4. Haplotypes Identified in the CTLA4 Gene

HAP No.	HAP ID	Polymorphic Sites						
		PS 1	PS 2	PS 3	PS 4	PS 5	PS 6	PS 7
1	745030	C	A	A	T	C	A	A
2	745025	C	A	G	T	C	A	A
3	745029	C	A	G	T	C	A	G
4	745031	C	A	G	T	C	G	A
5	745028	C	A	G	T	T	A	A
6	745032	C	G	G	C	C	A	A
7	745026	C	G	G	T	C	A	A
8	745027	T	A	G	T	C	A	A

Table 5 below shows the percent of chromosomes characterized by a given CTLA4 haplotype for all unrelated individuals in the Index Repository for which haplotype data was obtained. The percent of these unrelated individuals who have a given CTLA4 haplotype pair is shown in Table 6. In Tables 5 and 6, the "Total" column shows this frequency data for all of these unrelated individuals, while the other columns show the frequency data for these unrelated individuals categorized according to their self-identified ethnogeographic origin. Abbreviations used in Tables 5 and 6 are AF = African Descent, AS = Asian, CA = Caucasian, HL = Hispanic-Latino, and NA = Native American.

Table 5. Frequency of Observed CTLA4 Haplotypes In Unrelated Individuals

HAP No.	HAP ID	Total	CA	AF	AS	HL	NA
1	745030	0.61	0.0	2.5	0.0	0.0	0.0
2	745025	48.78	54.76	50.0	25.0	63.89	66.67
3	745029	0.61	2.38	0.0	0.0	0.0	0.0
4	745031	0.61	0.0	2.5	0.0	0.0	0.0
5	745028	0.61	0.0	2.5	0.0	0.0	0.0
6	745032	0.61	0.0	2.5	0.0	0.0	0.0
7	745026	42.07	38.1	40.0	60.0	30.56	33.33
8	745027	6.1	4.76	0.0	15.0	5.56	0.0

Table 6. Frequency of Observed CTLA4 Haplotype Pairs In Unrelated Individuals

HAP1	HAP2	Total	CA	AF	AS	HL	NA
2	2	23.17	23.81	25.0	5.0	38.89	33.33
7	7	19.51	14.29	25.0	35.0	5.56	0.0
2	1	1.22	0.0	5.0	0.0	0.0	0.0
2	3	1.22	4.76	0.0	0.0	0.0	0.0
2	5	1.22	0.0	5.0	0.0	0.0	0.0
2	6	1.22	0.0	5.0	0.0	0.0	0.0
2	8	6.1	9.52	0.0	10.0	5.56	0.0
2	4	1.22	0.0	5.0	0.0	0.0	0.0
7	8	6.1	0.0	0.0	20.0	5.56	0.0
2	7	39.02	47.62	30.0	30.0	44.44	66.67

The size and composition of the Index Repository were chosen to represent the genetic diversity across and within four major population groups comprising the general United States population. For example, as described in Table 1 above, this repository contains approximately equal sample sizes of African-descent, Asian-American, European-American, and Hispanic-Latino population groups. Almost all individuals representing each group had all four grandparents with the same ethnogeographic background. The number of unrelated individuals in the Index Repository provides a sample size that is sufficient to detect SNPs and haplotypes that occur in the general population with high statistical certainty. For instance, a haplotype that occurs with a frequency of 5% in the general population has a probability higher than 99.9% of being observed in a sample of 80 individuals from the general population. Similarly, a haplotype that occurs with a frequency of 10% in a specific population group has a 99% probability of being observed in a sample of 20 individuals from that population group. In addition, the size and composition of the Index Repository means that the relative frequencies determined therein for the haplotypes and haplotype pairs of the CTLA4 gene are likely to be similar to the relative frequencies of these CTLA4 haplotypes and haplotype pairs in the general U.S. population and in the four population groups represented in the Index Repository. The genetic diversity observed for the three Native Americans is presented because it is of scientific interest, but due to the small sample size it lacks statistical significance.

In view of the above, it will be seen that the several advantages of the invention are achieved and other advantageous results attained.

As various changes could be made in the above methods and compositions without departing from the scope of the invention, it is intended that all matter contained in the above description and shown in the accompanying drawings shall be interpreted as illustrative and not in a limiting sense.

All references cited in this specification, including patents and patent applications, are hereby incorporated in their entirety by reference. The discussion of references herein is intended merely to summarize the assertions made by their authors and no admission is made that any reference constitutes prior art. Applicants reserve the right to challenge the accuracy and pertinency of the cited references.

What is Claimed is:

1. A method for haplotyping the cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene of an individual, which comprises determining which of the CTLA4 haplotypes shown in the table immediately below defines one copy of the individual's CTLA4 gene, wherein each of the CTLA4 haplotypes comprises a set of polymorphisms whose locations and identities are set forth in the table immediately below:

Haplotype Number ^a								PS No. ^b	Nucleotide Position ^c	SEQ ID NO	Region Examined ^d
1	2	3	4	5	6	7	8				
C	C	C	C	C	C	C	T	1	37535	36	37286 - 37909
A	A	A	A	A	G	G	A	2	37902	36	37286 - 37909
A	G	G	G	G	G	G	G	3	38038	36	37608 - 38295
T	T	T	T	T	C	T	T	4	40867	36	40466 - 41010
C	C	C	C	T	C	C	C	5	41547	36	41123 - 41644
A	A	A	G	A	A	A	A	6	42460	36	42294 - 42918
A	A	G	A	A	A	A	A	7	42508	36	42294 - 42918

^aAlleles for haplotypes are presented 5' to 3' in each column

^bPS = polymorphic site;

^cLocation of PS within the indicated SEQ ID NO, wherein Nt = nucleotide;

^dRegion examined represents the nucleotide positions defining the start and stop positions of the sequenced region.

2. The method of claim 1, wherein the determining step comprises identifying the phased sequence of nucleotides present at each of PS1-7 on the one copy of the individual's CTLA4 gene.
3. A method for haplotyping the cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene of an individual, which comprises determining which of the CTLA4 haplotype pairs shown in the table immediately below defines both copies of the individual's CTLA4 gene, wherein each of the CTLA4 haplotype pairs consists of first and second haplotypes which comprise first and second sets of polymorphisms whose locations and identities are set forth in the table immediately below:

Haplotype Pairs ^a										PS No. ^b	Nucleotide Position ^c	SEQ ID NO	Region Examined ^d
2/2	7/7	2/1	2/3	2/5	2/6	2/8	2/4	7/8	2/7				
C/C	C/C	C/C	C/C	C/C	C/C	C/T	C/C	C/T	C/C	1	37535	36	37286 - 37909
A/A	G/G	A/A	A/A	A/A	A/G	A/A	A/A	G/A	A/G	2	37902	36	37286 - 37909
G/G	G/G	G/A	G/G	G/G	G/G	G/G	G/G	G/G	G/G	3	38038	36	37608 - 38295
T/T	T/T	T/T	T/T	T/T	T/C	T/T	T/T	T/T	T/T	4	40867	36	40466 - 41010
C/C	C/C	C/C	C/C	C/T	C/C	C/C	C/C	C/C	C/C	5	41547	36	41123 - 41644
A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/G	A/A	A/A	6	42460	36	42294 - 42918
A/A	A/A	A/A	A/G	A/A	A/A	A/A	A/A	A/A	A/A	7	42508	36	42294 - 42918

^aHaplotype pairs are represented as 1st Haplotype/2nd Haplotype; with alleles of each haplotype shown 5' to 3' as 1st Nt/2nd Nt in each column, where Nt = nucleotide;

^bPS = polymorphic site;

^cLocation of PS within the indicated SEQ ID NO, wherein Nt = nucleotide;

^dRegion examined represents the nucleotide positions defining the start and stop positions of

the sequenced region.

4. The method of claim 3, wherein the determining step comprises identifying the phased sequence of nucleotides present at each of PS1-7 on both copies of the individual's CTLA4 gene.
5. A method for genotyping the cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene of an individual, comprising determining for the two copies of the CTLA4 gene present in the individual the identity of the nucleotide pair at one or more polymorphic sites (PS) selected from the group consisting of PS1, PS3, PS4, PS5, PS6 and PS7, wherein the one or more PS have the location and alternative alleles shown in SEQ ID NO:36.
- 5 6. The method of claim 5, wherein the determining step comprises:
 - (a) isolating from the individual a nucleic acid mixture comprising both copies of the CTLA4 gene, or a fragment thereof, that are present in the individual;
 - (b) amplifying from the nucleic acid mixture a target region containing the selected polymorphic site;
 - 5 (c) hybridizing a primer extension oligonucleotide to one allele of the amplified target region;
 - (d) performing a nucleic acid template-dependent, primer extension reaction on the hybridized genotyping oligonucleotide in the presence of at least two different terminators of the reaction, wherein said terminators are complementary to the alternative nucleotides present at the selected polymorphic site; and
 - 10 (e) detecting the presence and identity of the terminator in the extended genotyping oligonucleotide.
7. The method of claim 5, which comprises determining for the two copies of the CTLA4 gene present in the individual the identity of the nucleotide pair at each of PS1-7.
8. A method for haplotyping the cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene of an individual which comprises determining, for one copy of the CTLA4 gene present in the individual, the identity of the nucleotide at two or more polymorphic sites (PS) selected from the group consisting of PS1, PS3, PS4, PS5, PS6 and PS7, wherein the two or more PS have the location and alternative alleles shown in SEQ ID NO:36.
9. The method of claim 8, further comprising determining the identity of the nucleotide at PS2, which has the location and alternative alleles shown in SEQ ID NO:36.
10. The method of claim 8, wherein the determining step comprises:
 - (a) isolating from the individual a nucleic acid sample containing only one of the two copies of the CTLA4 gene, or a fragment thereof, that is present in the individual;
 - (b) amplifying from the nucleic acid molecule a target region containing the selected polymorphic site;
 - 5 (c) hybridizing a primer extension oligonucleotide to one allele of the amplified target region;
 - (d) performing a nucleic acid template-dependent, primer extension reaction on the hybridized

genotyping oligonucleotide in the presence of at least two different terminators of the reaction, wherein said terminators are complementary to the alternative nucleotides present at the selected polymorphic site; and

- (e) detecting the presence and identity of the terminator in the extended genotyping oligonucleotide.

11. A method for predicting a haplotype pair for the cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene of an individual comprising:

- (a) identifying a CTLA4 genotype for the individual, wherein the genotype comprises the nucleotide pair at two or more polymorphic sites (PS) selected from the group consisting of PS1, PS3, PS4, PS5, PS6 and PS7, having the location and alternative alleles shown in SEQ ID NO:36;
- (b) enumerating all possible haplotype pairs which are consistent with the genotype;
- (c) comparing the possible haplotype pairs to the haplotype pair data set forth in the table immediately below; and
- (d) assigning a haplotype pair to the individual that is consistent with the data

Haplotype Pairs ^a										PS No. ^b	Nucleotide Position ^c	SEQ ID NO	Region Examined ^d
2/2	7/7	2/1	2/3	2/5	2/6	2/8	2/4	7/8	2/7				
C/C	C/C	C/C	C/C	C/C	C/C	C/T	C/C	C/T	C/C	1	37535	36	37286 - 37909
A/A	G/G	A/A	A/A	A/A	A/G	A/A	A/A	G/A	A/G	2	37902	36	37286 - 37909
G/G	G/G	G/A	G/G	G/G	G/G	G/G	G/G	G/G	G/G	3	38038	36	37608 - 38295
T/T	T/T	T/T	T/T	T/T	T/C	T/T	T/T	T/T	T/T	4	40867	36	40466 - 41010
C/C	C/C	C/C	C/C	C/T	C/C	C/C	C/C	C/C	C/C	5	41547	36	41123 - 41644
A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/G	A/A	A/A	6	42460	36	42294 - 42918
A/A	A/A	A/A	A/G	A/A	A/A	A/A	A/A	A/A	A/A	7	42508	36	42294 - 42918

^aHaplotype pairs are represented as 1st Haplotype/2nd Haplotype; with alleles of each haplotype shown 5' to 3' as 1st Nt/2nd Nt in each column, where Nt = nucleotide;

^bPS = polymorphic site;

^cLocation of PS within the indicated SEQ ID NO, wherein Nt = nucleotide;

^dRegion examined represents the nucleotide positions defining the start and stop positions of the sequenced region.

12. The method of claim 11, wherein the identified genotype of the individual comprises the nucleotide pair at each of PS1-7, which have the location and alternative alleles shown in SEQ ID NO:36.
13. A method for identifying an association between a trait and at least one haplotype or haplotype pair of the cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene which comprises comparing the frequency of the haplotype or haplotype pair in a population exhibiting the trait with the frequency of the haplotype or haplotype pair in a reference population, wherein the haplotype is selected from haplotypes 1-8 shown in the table presented immediately below, wherein each of the haplotypes comprises a set of polymorphisms whose locations and identities are set forth in the table immediately below:

Haplotype Number ^a								PS No. ^b	Nucleotide Position ^c	SEQ ID NO	Region Examined ^d
1	2	3	4	5	6	7	8				
C	C	C	C	C	C	C	T	1	37535	36	37286 - 37909
A	A	A	A	A	G	G	A	2	37902	36	37286 - 37909
A	G	G	G	G	G	G	G	3	38038	36	37608 - 38295
T	T	T	T	T	C	T	T	4	40867	36	40466 - 41010
C	C	C	C	T	C	C	C	5	41547	36	41123 - 41644
A	A	A	G	A	A	A	A	6	42460	36	42294 - 42918
A	A	G	A	A	A	A	A	7	42508	36	42294 - 42918

^aAlleles for haplotypes are presented 5' to 3' in each column

^bPS = polymorphic site;

^cLocation of PS within the indicated SEQ ID NO, wherein Nt = nucleotide;

^dRegion examined represents the nucleotide positions defining the start and stop positions of the sequenced region,

and the haplotype pair is selected from the haplotype pairs shown in the table immediately below, wherein each of the CTLA4 haplotype pairs consists of first and second haplotypes which comprise first and second sets of polymorphisms whose locations and identities are set forth in the table immediately below:

Haplotype Pairs ^a										PS No. ^b	Nucleotide Position ^c	SEQ ID NO	Region Examined ^d
2/2	7/7	2/1	2/3	2/5	2/6	2/8	2/4	7/8	2/7				
C/C	C/C	C/C	C/C	C/C	C/C	C/T	C/C	C/T	C/C	1	37535	36	37286 - 37909
A/A	G/G	A/A	A/A	A/A	A/G	A/A	A/A	G/A	A/G	2	37902	36	37286 - 37909
G/G	G/G	G/A	G/G	G/G	G/G	G/G	G/G	G/G	G/G	3	38038	36	37608 - 38295
T/T	T/T	T/T	T/T	T/T	T/C	T/T	T/T	T/T	T/T	4	40867	36	40466 - 41010
C/C	C/C	C/C	C/C	C/T	C/C	C/C	C/C	C/C	C/C	5	41547	36	41123 - 41644
A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/G	A/A	A/A	6	42460	36	42294 - 42918
A/A	A/A	A/A	A/G	A/A	A/A	A/A	A/A	A/A	A/A	7	42508	36	42294 - 42918

^aHaplotype pairs are represented as 1st Haplotype/2nd Haplotype; with alleles of each haplotype shown 5' to 3' as 1st Nt/2nd Nt in each column, where Nt = nucleotide;

^bPS = polymorphic site;

^cLocation of PS within the indicated SEQ ID NO, wherein Nt = nucleotide;

^dRegion examined represents the nucleotide positions defining the start and stop positions of the sequenced region,

wherein a higher frequency of the haplotype or haplotype pair in the trait population than in the reference population indicates the trait is associated with the haplotype or haplotype pair.

14. The method of claim 13, wherein the trait is a clinical response to a drug targeting CTLA4.
15. A composition comprising at least one genotyping oligonucleotide for detecting a polymorphism in the cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene at a polymorphic site (PS) selected from the group consisting of PS1, PS3, PS4, PS5, PS6 and PS7, having the location and alternative alleles shown in SEQ ID NO:36.
16. The composition of claim 15, wherein the genotyping oligonucleotide is an allele-specific oligonucleotide that specifically hybridizes to an allele of the CTLA4 gene at a region containing

- the polymorphic site.
17. The composition of claim 16, wherein the allele-specific oligonucleotide comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS:4-9, the complements of SEQ ID NOS:4-9, and SEQ ID NOS:10-21.
 18. The composition of claim 15, wherein the genotyping oligonucleotide is a primer-extension oligonucleotide.
 19. The composition of claim 18, wherein the primer extension oligonucleotide comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS:22-33.
 20. A kit for genotyping the cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene of an individual, which comprises a set of oligonucleotides designed to genotype each of polymorphic sites (PS) PS1, PS3, PS4, PS5, PS6 and PS7, having the location and alternative alleles shown in SEQ ID NO:36.
 21. The kit of claim 20, which further comprises oligonucleotides designed to genotype PS2, having the location and alternative alleles shown in SEQ ID NO:36.
 22. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
 - (a) a first nucleotide sequence which comprises a cytotoxic T-lymphocyte-associated protein 4 (CTLA4) isogene, wherein the CTLA4 isogene is selected from the group consisting of isogenes 2-8 shown in the table immediately below and wherein each of the isogenes comprises the regions of the SEQ ID NOS shown in the table immediately below and wherein each of the isogenes 2-8 is further defined by the corresponding set of polymorphisms whose locations and polymorphisms are set forth in the table immediately below

Isogene Number ^a								PS No. ^b	Nucleotide Position ^c	SEQ ID NO	Region Examined ^d
1	2	3	4	5	6	7	8				
C	C	C	C	C	C	C	T	1	37535	36	37286 - 37909
A	A	A	A	A	G	G	A	2	37902	36	37286 - 37909
A	G	G	G	G	G	G	G	3	38038	36	37608 - 38295
T	T	T	T	T	C	T	T	4	40867	36	40468 - 41010
C	C	C	C	T	C	C	C	5	41547	36	41123 - 41644
A	A	A	G	A	A	A	A	6	42460	36	42294 - 42918
A	A	G	A	A	A	A	A	7	42508	36	42294 - 42918

^aAlleles for isogenes are presented 5' to 3' in each column

^bPS = polymorphic site;

^cLocation of PS within the indicated SEQ ID NO, wherein Nt = nucleotide;

^dRegion examined represents the nucleotide positions defining the start and stop positions of the sequenced region;

- (b) a second nucleotide sequence which comprises a fragment of the first nucleotide sequence, wherein the fragment comprises one or more polymorphisms selected from the group consisting of thymine at PS1, guanine at PS3, cytosine at PS4, thymine at PS5, guanine at

PS6 and guanine at PS7, wherein the selected polymorphism has the location set forth in the table immediately above; and

(c) a third nucleotide sequence which is complementary to the first or second nucleotide sequence.

23. The isolated polynucleotide of claim 22, which is a DNA molecule and comprises both the first and third nucleotide sequences and further comprises expression regulatory elements operably linked to the first nucleotide sequence.
24. A recombinant nonhuman organism transformed or transfected with the isolated polynucleotide of claim 22, wherein the organism expresses a CTLA4 protein encoded by the first nucleotide sequence.
25. The recombinant organism of claim 24, which is a nonhuman transgenic animal.
26. The isolated polynucleotide of claim 22 which consists of the second nucleotide sequence.
27. A computer system for storing and analyzing polymorphism data for the cytotoxic T-lymphocyte-associated protein 4 gene, comprising:
 - (a) a central processing unit (CPU);
 - (b) a communication interface;
 - (c) a display device;
 - (d) an input device; and
 - (e) a database containing the polymorphism data;

wherein the polymorphism data comprises the haplotypes set forth in the table immediately below:

Haplotype Number ^a								PS No. ^b	Nucleotide Position ^c	SEQ ID NO	Region Examined ^d
1	2	3	4	5	6	7	8				
C	C	C	C	C	C	C	T	1	37535	36	37286 - 37909
A	A	A	A	A	G	G	A	2	37902	36	37286 - 37909
A	G	G	G	G	G	G	G	3	38038	36	37608 - 38295
T	T	T	T	T	C	T	T	4	40867	36	40466 - 41010
C	C	C	C	T	C	C	C	5	41547	36	41123 - 41644
A	A	A	G	A	A	A	A	6	42460	36	42294 - 42918
A	A	G	A	A	A	A	A	7	42508	36	42294 - 42918

^aAlleles for haplotypes are presented 5' to 3' in each column

^bPS = polymorphic site;

^cLocation of PS within the indicated SEQ ID NO, wherein Nt = nucleotide;

^dRegion examined represents the nucleotide positions defining the start and stop positions of the sequenced region;

and the haplotype pairs set forth in the table immediately below:

Haplotype Pairs ^a										PS No. ^b	Nucleotide Position ^c	SEQ ID NO	Region Examined ^d
2/2	7/7	2/1	2/3	2/5	2/6	2/8	2/4	7/8	2/7				
C/C	C/C	C/C	C/C	C/C	C/C	C/T	C/C	C/T	C/C	1	37535	36	37286 - 37909
A/A	G/G	A/A	A/A	A/A	A/G	A/A	A/A	G/A	A/G	2	37902	36	37286 - 37909
G/G	G/G	G/A	G/G	G/G	G/G	G/G	G/G	G/G	G/G	3	38038	36	37608 - 38295
T/T	T/T	T/T	T/T	T/T	T/C	T/T	T/T	T/T	T/T	4	40867	36	40466 - 41010
C/C	C/C	C/C	C/C	C/T	C/C	C/C	C/C	C/C	C/C	5	41547	36	41123 - 41644
A/A	A/A	A/A	A/A	A/A	A/A	A/A	A/G	A/A	A/A	6	42460	36	42294 - 42918
A/A	A/A	A/A	A/G	A/A	A/A	A/A	A/A	A/A	A/A	7	42508	36	42294 - 42918

^aHaplotype pairs are represented as 1st Haplotype/2nd Haplotype; with alleles of each haplotype shown 5' to 3' as 1st Nt/2nd Nt in each column, where Nt = nucleotide;

^bPS = polymorphic site;

^cLocation of PS within the indicated SEQ ID NO, wherein Nt = nucleotide;

^dRegion examined represents the nucleotide positions defining the start and stop positions of the sequenced region.

28. A genome anthology for the cytotoxic T-lymphocyte-associated protein 4 (CTLA4) gene which comprises CTLA4 isogenes defined by any one of haplotypes 1-8 set forth in the table shown below:

Haplotype Number ^a								PS No. ^b	Nucleotide Position ^c	SEQ ID NO	Region Examined ^d
1	2	3	4	5	6	7	8				
C	C	C	C	C	C	C	T	1	37535	38	37286 - 37909
A	A	A	A	A	G	G	A	2	37902	38	37286 - 37909
A	G	G	G	G	G	G	G	3	38038	38	37608 - 38295
T	T	T	T	T	C	T	T	4	40867	38	40466 - 41010
C	C	C	C	T	C	C	C	5	41547	38	41123 - 41644
A	A	A	G	A	A	A	A	6	42460	38	42294 - 42918
A	A	G	A	A	A	A	A	7	42508	38	42294 - 42918

^aAlleles for haplotypes are presented 5' to 3' in each column

^bPS = polymorphic site;

^cLocation of PS within the indicated SEQ ID NO, wherein Nt = nucleotide;

^dRegion examined represents the nucleotide positions defining the start and stop positions of the sequenced region.

1/5

POLYMORPHISMS IN THE CTLA4 GENE

ACTCAAATTT	CCCCCAGTTT	CTCAGTAATC	TGCTTTGTAT	TATTTTTTTC	
AATCAATAAT	AACACATTAA	ATTCTGTCAT	CATGTCTCTA	TTTTCTCCTT	33900
TAATCTAGAA	TGGTTTTCCA	GGCTTTTCCC	TTTTTAATCT	TTTGTGATGT	
TAACATTTTT	GAAGAATCTA	GGCCAGTTAT	TTTGCAGAAT	ATGGATTTGT	34000
CTGATTATTT	CTTCATGCTT	AGATTCAGTT	AAAACAATTT	TGGCAATAAC	
TACACAAAGT	TGTTGTGTCC	TTCTCAGTGT	ATTGTATCTG	GAGGTACCCT	34100
ATGTCAGAAC	ACTCCATCAC	AGTAATCTTC	AGTTGGGATC	ACTTGGTTAA	
AGCAGTATCT	GCTAGATTTA	TCAATTATCT	TTTCCCTATT	GTAATTATCA	34200
AGTAATCTGT	GGGGTGGTAT	ATGTGAATAT	ATCTTTTCTC	ACCAAGATTTC	
TTTTACCCAG	CATTTTTTTT	ACAACCATT	GAATTTTCTT	GCTATGACAG	34300
AATTAGTTTC	ATAAGAGCAT	AATTAAGGAA	TCTGGTAAGA	CGTCTTTGAA	
AGTTCCTAAT	ACAACCCTCA	TCTCTTCTTA	TGTTAAACTG	TCTTCATTCA	34400
TTTACCTACC	TATTTAACAA	ATGTTTAAAG	AATGCTCGTT	TTGTGGCTTG	
GCCTGTGACA	TTTTCATCAG	TGGCTTCATA	GTTGCCTAAA	CTAATGTCTC	34500
TTCTGTGCT	CCAGAATCCT	ATTGGCAACT	ATTTTCCATG	GATGCCATAA	
ACTCAACATA	CTCAGATGAA	AGCTGTCAAT	TCACCTTCTC	ACCTGCTCTT	34600
CCTGTTATTT	ATTCCAGTAA	ATTGCACCTC	TGCTTCTCTA	TTGACCCTAT	
AATAAGAAAG	GCCTATTTAT	TGAGTGGCTA	TCTGTTACAA	TAGGCCTTTT	34700
ATACATGTTA	CATGCATTAT	CTCATTTAAT	CTTTTCAATA	ATCTTGAGAA	
TTAGATGCCA	TTATCTGGAG	GACATTGGGC	TTTGAGCCCC	AGGTCAACAG	34800
TAAGCAAGTC	CCCATGCTAG	GATACAAATC	CAGTTATCTT	GGGCTTCAAA	
ATCTTTGCTT	TAGCATAGTG	ATAAAGGTAT	CACTGGCTTC	AAATCCTGGC	34900
TCTGCCACTT	CCTACGGGTG	ATTGGGTAAG	TCACCTTAAC	TTTCAATGCC	
TTGATTTTCT	TCTTTATAAA	ATGGGAAAAA	TGGTAACTCT	TGTCTTGTA	35000
GGTTGTTATG	GACTTGAAAG	AAACAAAAC	TGCTGGGCGC	GGTGGCTCAC	
GCCTGTAATC	CCAGCGCTTT	GGGAGGCCGA	GGCGGGTAGA	TCACCTGAGG	35100
TCAGGAATTT	GAGACTAGCC	TGGTCAACAT	GGCGAAACCC	CATCTCTACT	
AAAAATAATC	AAAAAATTAG	CTGGGCGTGG	TGGTGGGTGC	CTGCAATCCC	35200
AGCTACTTGG	GAGGCTGAGG	CAGGAGAATC	ACTTGAACCC	AGGAGGCAGA	
GGTTGCAGTG	AGCCGAGATC	GCGCCACTGC	ACTCCAACCT	GGGAAACAAG	35300
AGTGAAACTC	TGTCTCAAAA	AAATAAAAAA	GAAAAGAAAG	AAAACCTGAC	
ATTTGAATAA	TTTCTGCAGC	ACCACAGTAC	TTCTCAAACC	ATAATTCTGG	35400
TGTCATTCAT	TCTAAACTTC	TCCTTCCACT	CACTGATTGT	GGACTAGTCC	
CATAGGGACC	TCGTCTTCAC	AGCTGTCAAT	ACCTTAGTTT	AGACCAGGGG	35500
TTGGCAAACC	ATGGCCCACA	GGCCAAATCC	AGACTTCACC	TATTTTCGTA	
AATGAAGTTT	TATTGGAATG	CACACACACC	CATTGTGTTA	CATATAATAT	35600
ATGGCTTCTT	TCATTCTAAA	ATGACAGAGT	TGAGTAGTGG	CAACAGAGAC	
CCCACCGTTT	GCAAATCATA	ACATATTTAC	TATTTTGCCC	CCTTCAGAAA	35700
GCTTTCCAAA	TCCTGGTGCC	TGGCGTATTC	CAATAGTCTT	CTTCCCAGTC	
TTCTGGGTTA	CATTTCTCCC	TGAACCCATC	CTCCCCATCC	CTAAAAATTC	35800
CCCCAAATGT	GAACCTAGTC	ATACCAGTTT	GCTCCTTATG	TTTCATTGGC	
CCTTGCTGCT	AAGAGCATCC	GCTTGCACCT	TCTGCTCATC	CCCAGACAAG	35900
CTTTGTCTCG	TGACCATAAT	GAACCTTTCA	TGCCGTTTCC	AACTTTAGCC	
CATGTTATTC	TTCTTGCTCG	AATATCCACC	CTTTTCTCTG	TTCTCAATAA	36000
TAAGTTCAGG	CTTTTCTGCT	TCTGAGAAGC	CCTTTCTGAC	TTCCACAGGC	
TGAACCACTG	GCTTCTGCTC	CTCTACATAA	TACTTCAATT	CCAGCATTGA	36100
TCTCACTCTA	TCATGATCAT	GGGTTTAGCT	GTCTGTCCCT	GCCACTGCTG	
TGTGTTCCCT	TTGAGGGCAG	GAACATTTGT	TTTTCACTTT	TTAAAAAACC	36200
TCTGTTGCCC	AGTCTGGCAT	TAGGAAGTGC	CCATTAGGTT	GTTATTGCTT	
GTTGGCGCTT	GAGCTGGGGC	TTGAAGGTTT	CTATAATGTG	TAGCAGTGTA	36300
TAGAAAACAG	GCAGGTCAGA	AAAGGCTTCT	GTGCATCACA	CCAACATGGC	

FIGURE 1A

2/5

ACRTGTATAC ATATGTAACA AATCTGCATG TTGTGCACAT GTACCCTAAA 36400
 ACTTAAAGTA TAATAATAAT AAAATTTTAA AAAAAAAG AAGAGGCTTC
 CTGGAGGAGA TGACAGCTGA GCTAAGTCCT GGAGGATGAG AAGGAGTATA 36500
 AAATAAGATA ATAGGAGAAA AAAGGCAGTA GGAACAGCAT GGGTAAAGGT
 GATGAGGCCT GAAAGAGGCCA CGTGGAAAGGA AAGACAAATG CAGGAAGGGG 36600
 GAATGGGAGG GAATGCTGGG GTACAGGCCA AAGAGGGAGG CATTGCGGGA
 GTGTTCTTTT AAACATCCTT TGGTGTGCTA ATGTGTGGTC ATTGGGAAAC 36700
 CATGGACCGT TTTTTTTTGG CAAACCTCAT ATCCCCTGTT ACAACTGTCT
 GTTTGCATGT CAGCCTTGTA AAAGCCCCTT AAGGTATCAA CTATGTTTTT 36800
 GTTTTGTGAT CATTCAATCC TAAGTGCACA GAATCCCGG CATATTACAG
 GTTCCCATG AATGTTTCTT TCTTTATTAA AATGTATGAA AACTCTCCAG 36900
 ATTTAAGGAA GGTCCTCAAT GTTTCAAATT CTTTTGTTA GATCATTGGT
 CCTGTCTACA GCTGTACAAA ATTTAAGGAC TCTGTTTATA TTTAATCTTC 37000
 ACTTTTGAAT TTTCTGCTTG AAAAATTTGT ATTAGAAAAA AAAGTCTATC
 CTTTTATGGA CGGCTCTAAT CTCTTGAATC ATTTGGGTTG GCTTTTCTTT 37100
 GGACCTTCTT CAACTCTGTT TTGTCTCTGT TGAGTTAAGG CTTTTAAGAA
 CACCTGAATT CTTTCCTTCT GCAAAACCAG AGGCAGCTTC TTTCCGCCT 37200
 ATTTTCAGTT TATTTCTTGT GATTTTAGTT TTTTCTCTT AACCAAATGC
 TAAATGGATT TAGGAGAAAT AAACCTATTT GTAAAGCTGT CAAGGGACCA 37300
 TTAGAAGGAT GGTGCTTCAC AGATAGAATA CAGTTTTTAT TAATGATGCC
 TAGACAAATC CTGCCATTAG CCCAAGGGCT CAGAAAGTTA GCAGCCTAGT 37400
 AGTTTGGAG TTGTCAATGA AATGAATTGG ACTGGATGGT TAAGGATGCC
 CAGAGATTG AATAAAATTG GGATTTAGGA GGACCCTTGT ACTCCAGGAA 37500
 ATTCTCCAAG TCTCCACTTA GTTATCCAGA TCCTCAAAGT GAACATGAAG
 T
 CTTCACTTTC AAATTGAATA CATTTTCCAT CCATGGATTG GCTTGTTTTG 37600
 TTCAGTTGAG TGCTTGAGGT TGTCTTTTCG ACGTAACAGC TAAACCCACG
 GCTTCTTTTC TCGTAAAACC AAAACAAAAA GGCTTTCTAT TCAAGTGCTT 37700
 TCTGTGTGTG CACATGTGTA ATACATATCT GGGATCAAAG CTATCTATAT
 AAAGTCCTTG ATTCTGTGTG GGTTCAAACA CATTTCAAAG CTTCAGGATC 37800
 CTGAAAGGTT TTGCTCTACT TCCTGAAGAC CTGAACACCG CTCCCATAAA
 GCCATGGCTT GCCTTGGATT TCAGCGGCAC AAGGCTCAGC TGAACCTGGC 37900
 [exon 1: 37854..
 TACCAGGACC TGGCCCTGCA CTCTCCTGTT TTTTCTTCTC TTCATCCCTG
 G
 TCTTCTGCAA AGGTGAGTGA GACTTTTGGG GCATGAAGAT GGAGGAGGTG 38000
 ...37961]
 TTTCTCTTAC CTGGGTTTCA TTTGTTTCAG CAGTCAAAGG CAGTGATTTA
 G
 TAGCAAAGCC AGAAGTTAAA GGTAAACTC CAATCTGGCT TGGCTGGCTC 38100
 TGTATCCAG GGCCAGCAGG GAGCAGTTGG GCGGCACGAA ATAAGGCAAA
 GAGATAGCTC AGAACAGAGC GCCAGGTATT TAGTAGGGGC TTCATGAATG 38200
 CATGTGAGTT GGTTAGTAG AGAGACACAG GCAATTTTCAG ACCCTTCTAT
 GAGATGGGAA GTGATTTAAG AGGGAAAGGA TAGCCATAGT CCTGAATACA 38300
 TTTGAGCTGG GTTTCAGGAT GAGCTCACAA GTTCCTTTAA AAAAAATTGA
 CTTAAGCAAA TCCTGGGAAG AGTTTTTTTG CTATACAATT CAAGGTTTTA 38400
 AGGTCCTCGG ATTCATATAC TTTATAAATG AATTAGCCAG CTTGTTTTAA
 ATGTAGGGAA ATTGTGGGAA GAATGCCTTC TTTACTTAAT TCAAGGTTTT 38500
 AAGGTTCTCT TAATCAATTC TACTAGCTAA TTAGCCAATT ATTTAAAAAT
 AAAAGTTTGA AATTGCCAAA AAAAAAGAC AAGGAAAAGG AAAGAAAGAA 38600
 AGCCACCAGT CTGTTTGGCA TACAATACTT AATTGTTGCC TGACCTACGT
 GTGGGTTTCA GATGCAGATC CTCAGTTTTC AGCTCTTCAG AGACTGACAC 38700
 CAGGTTTGTG ACACGGCTTA AAATGATGAG TATATCCATT GAATCTCAAC

FIGURE 1B

3/6

CTTATCTCTC	TCTAGACCTT	CTTGGTTAAG	AAACCATGTA	GTTTGTATGA	38800
AGTAGGTACT	CAAAAAGATAT	TTGATGATTT	AATTTTACT	GGAGAAGAAA	
TATTCATATA	TGTTTTCTTA	TTTTTACATG	TTTTAAATAT	GTAAAGATTA	38900
AATAAACACT	CTTAGAAGTA	TTTAAATTC	CTAAAGTAAA	TTTATCTCAA	
CCAGTAACAG	GACCCTCCCA	ATACTGGAAA	GTTGAGTGTG	ACCGCATTTA	39000
TGGTGATGA	GTGTGAGCTT	GCTTGGGGAG	AGGGCAGGAC	ATTTAGGATT	
TCTTAAGCTT	AGAGTCAATA	CAATAAAGAT	TATTGAGTGC	TCACTGGGT	39100
GGGCTATAAT	CACTGCTCAC	AGGAGTTCAT	GAACCACAAG	TAAAAGAGTG	
AGGAGATATG	ATTAGCTCAC	AAATAACTTT	AATACAGAGC	AGAAAGTAAT	39200
GAAGTACTGC	AATGGAGTTA	TCACAGTGCT	AAGGATGCTC	AGAGGGCATC	
TCTGATAGGC	AGAGGTGAGG	GTTAGGGAAG	GAAGCTGTAG	TCTAGCTAGC	39300
TAGAGCTGCT	GGAATAGACA	TGACAATGGC	TGCTGCAAAC	TGTTTTCTCT	
TCTGAGGACA	GATGTCCCGT	GCAAGTGGCT	TGGTGGAAGG	GACTAGTGTG	39400
TCTAATATAG	GGTGATTTAT	AAGCAGGAAA	GTGTGTCCTA	GAAATTCAGA	
CCAGAGTGAT	AGATTGGAAT	TGGATCATGG	GGGACTCATT	GAATGTTATT	39500
TATTGTATTT	GTTTTTGCGA	TCAGTGTTAG	TAAAGTGTCA	AAGGGATTGA	
GCAGATGAGT	GACATCATGC	AACACAAGTT	TTGAGTTTCA	CTTGTGAGAC	39600
TGACTGGAGA	GGGGCCTGGT	TAGTTACAGG	AAGGTAATTT	GGCATGCAGC	
CACTATTTTT	GAGTTGATGC	AAGCCTCTCT	GTATGGAGAG	CTGGTCTCCT	39700
TTATCCTGTG	GGAAAAGAGA	ACAAAGGAGC	ATGGGAGTGT	TCAAGGGAAG	
GAGAAATAAA	GGGCAGAGAG	GCAGCGGTGG	TGTCAGGGGA	AGCCCACAGG	39800
AGTTAACAGC	AGGGTTGCCT	CAACCTAGAG	AGGAAGCGAC	CTGGTGCCCT	
CGGCTCTGTG	GCTTCCTTCA	TCTAACAACA	TCTTCCACTC	TACAACAATG	39900
CCAGGGAAGG	CGGAGGCTGG	TACAGTGCAT	CAAGACACAG	CTACTCCTGG	
GTGACAGAGG	TTCAGGGCCA	GCTCACTAAG	TAGGCAGAAG	TTTTTGACAT	40000
ATACTTTGAG	AGATAAAGCA	AGATTCTGTA	CCTCAACCTT	CAGAATTTCC	
CCTACCACTC	ATTATAGTTC	CGGAGCTATA	TAGCTCCTAT	CATTCTATCA	40100
TAACCTTAGA	ATACCAGAGA	ACATATCATC	TCATCTAATT	ATCTCTTACT	
ATATGTGAAA	AAAATGAAGG	ACATGGGGGA	AGTGTGACTT	GCCCCAAATC	40200
ACATATTTCA	TGGTAGAGCC	AGGTCTTCTG	TTTGTATAT	CAGTGTCTT	
CCTGCCACAA	CCATCTTGAA	GAATCTATTT	CTCAGTAAGA	AAATATCTTT	40300
ATGGAGAGTA	GCTGGAAAAC	AGTTGAGAGA	TGGAGGGGAG	GCTGGGGGTG	
TGGAGAGGGG	AAGGGGTAAAG	TGATAGATTC	GTTGAAGGGG	GGAGAAAAGG	40400
CCGTGGGGAT	GAAGCTAGAA	GGCAGAAGGG	CTTGCCCTGGG	CTTGGCCATG	
AAGGAGCATG	AGTTCATGA	GTTCCCTTTG	GCTTTTCCAT	GCTAGCAATG	40500
[exon 2: 40495..					
CACGTGGCCC	AGCCTGCTGT	GGTACTGGCC	AGCAGCCGAG	GCATCGCCAG	
CTTTGTGTGT	GAGTATGCAT	CTCCAGGCAA	AGCCACTGAG	GTCCGGGTGA	40600
CAGTGCTTCG	GCAGGCTGAC	AGCCAGGTGA	CTGAAGTCTG	TGCGGCAACC	
TACATGATGG	GGAATGAGTT	GACCTTCCTA	GATGATTCCA	TCTGCACGGG	40700
CACCTCCAGT	GGAAATCAAG	TGAACCTCAC	TATCCAAGGA	CTGAGGGCCA	
TGGACACGGG	ACTCTACATC	TGCAAGGTGG	AGCTCATGTA	CCCACCGCCA	40800
TACTACCTGG	GCATAGGCAA	CGGAACCCAG	ATTTATGTAA	TTGGTGAGCA	
..40843]					
AAGCCATTTT	ACTGAGTTGA	CACCTGTTGC	ATTGCAGTCT	TCTATGCACA	40900
C					
AAAACAGTTT	TGTTCTTTAA	TTTCAGGAGG	TTTACTTTTA	GGAAGTGGGA	
CATTCTCTTT	AAGAGTTCTG	TACCACATGG	TAGCCTTGCT	TATTGTGGGT	41000
GGCAACCTTA	ATAGCATTCT	GAAGTGTAAA	TAAAATGATT	TGGGGAAGTT	
GGGGCTCTCG	CTCTGGAGTG	CTAACCATCA	TGACGTTTGA	TCTGTACTTT	41100
TGATATGATA	TGATGCTCCT	GGGGAAGTAG	TCCCAAATAG	CCAAACCTAT	
TGGTGGGCTA	CCCATGCAAT	TTAGGGGTGG	ACCTCAAGGC	CTGGAAGCTC	41200
TAATGTCCTT	TTTTACCAA	TGTTGGGGAG	TAGAGCCCTA	GAGTTTAAAA	

FIGURE 1C

5/6

POLYMORPHISMS IN THE CODING SEQUENCE OF CTLA4

ATGGCTTGCC	TTGGATTTC	GCGGCACAAG	GCTCAGCTGA	ACCTGGCTAC	
				G	
CAGGACCTGG	CCCTGCACTC	TCCTGTTTTT	TCTTCTCTTC	ATCCCTGTCT	100
TCTGCAAAGC	AATGCACGTG	GCCCAGCCTG	CTGTGGTACT	GGCCAGCAGC	
CGAGGCATCG	CCAGCTTTGT	GTGTGAGTAT	GCATCTCCAG	GCAAAGCCAC	200
TGAGGTCCGG	GTGACAGTGC	TTCGGCAGGC	TGACAGCCAG	GTGACTGAAG	
TCTGTGCGGC	AACCTACATG	ATGGGGAATG	AGTTGACCTT	CCTAGATGAT	300
TCCATCTGCA	CGGGCACCTC	CAGTGGAAAT	CAAGTGAACC	TCACTATCCA	
AGGACTGAGG	GCCATGGACA	CGGGACTCTA	CATCTGCAAG	GTGGAGCTCA	400
TGTACCCACC	GCCATACTAC	CTGGGCATAG	GCAACGGAAC	CCAGATTTAT	
GTAATTGATC	CAGAACCGTG	CCCAGATTCT	GACTTCCTCC	TCTGGATCCT	500
TGCAGCAGTT	AGTTCGGGGT	TGTTTTTTTA	TAGCTTTCTC	CTCACAGCTG	
TTTCTTTGAG	CAAAATGCTA	AAGAAAAGAA	GCCCTCTTAC	AACAGGGGTC	600
TATGTGAAAA	TGCCCCCAAC	AGAGCCAGAA	TGTGAAAAGC	AATTCAGCC	
TTATTTTATT	CCCATCAATT	GA			672

FIGURE 2

SEQUENCE LISTING

<110> Genaiissance Pharmaceuticals, Inc.

Chew, Anne

Choi, Julie Y.

Messer, Chad

<120> Haplotypes of the CTLA4 Gene

<130> MWH-0537PCT CTLA4

<140> TBA

<141> 2001-05-23

<150> 60/206,353

<151> 2000-05-23

<160> 36

<170> PatentIn Ver. 2.1

<210> 1

<211> 9336

<212> DNA

<213> Homo sapien

<400> 1

```
actcaaattt cccccagttt ctcagtaatc tgctttgtat tatttttttc aatcaataat 60
aacacattaa attctgtcat catgtctcta ttttctcctt taatctagaa tggttttcca 120
ggcttttccc tttttaatct tttgtgatgt taacattttt gaagaatcta ggccagttat 180
tttgcagaat atggatttgt ctgattttt cttcatgctt agattcagtt aaaacaattt 240
tggcaataac tacacaaagt tgttgtgtcc ttctcagtgt attgtatctg gaggtaccct 300
atgtcagaac actccatcac agtaatcttc agttgggac acttgggtta agcagtatct 360
gctagattta tcaattatct tttccctatt gtaattatca agtaatctgt ggggtgggtat 420
atgtgaatat atcttttctc accaagattc ttttaccag catttttttt acaaccattt 480
gaattttctt gctatgacag aattagtttc ataagagcat aattaaggaa tctggttaaga 540
cgtctttgaa agttctaatt acaaccctca tctctcttta tgttaaactg tcttcattca 600
tttacctacc tatttaacaa atgtttaaag aatgctcgtt ttgtggcttg gcctgtgaca 660
ttttcatcag tggcttcata gttgcctaaa ctaatgtctc ttctgtgtct ccagaatcct 720
attggcaact attttccatg gatgccataa actcaacata ctcagatgaa agctgtcatt 780
tcactttctc acctgctctt cctgttattt attccagtaa attgcacctc tgtcttctca 840
ttgacctat aataagaaag gcctatttat tgagtggcta tctgttacia taggcctttt 900
atacatgtta catgcattat ctcatttaat cttttcaata atcttgagaa ttagatgcca 960
ttatctggag gacattgggc tttgagcccc aggtcaacag taagcaagtc ccatgctag 1020
gatacaaatc cagttatctt gggcttcaaa atctttgctt tagcatagtg ataaagggtat 1080
cactggcttc aaatcctggc tctgccactt cctacgggtg attgggtaag tcaccttaac 1140
tttcaatgcc ttgatttcct tctttataaa atgggaaaaa tggtaactct tgtctttag 1200
ggttgttatg gacttgaaag aaacaaaact tgctgggcgc ggtggctcac gcctgtaatc 1260
```

ccagcgctttt gggaggccga ggcgggtaga tcacctgagg tcaggaaatt gagactagcc 1320
tggccaacat ggcgaaaccc catctctact aaaaataaac aaaaaattag ctgggcgtgg 1380
tgggggtgc ctgcaatccc agctacttgg gaggttgagg caggagaatc acttgaaccc 1440
aggaggcaga ggttgcaagt agccgagatc gcgccactgc actccaacct gggaaacaag 1500
agtgaacctc tgtctcaaaa aaataaaaaa gaaaagaaaag aaaacttgac atttgaataa 1560
ttctgcagc accacagtac ttctcaaacc ataattctgg tgcattcat tctaaacttc 1620
tccttccact cactgattgt ggactagtcc catagggacc tegtcttcac agctgtcatt 1680
accttagtct agaccagggg ttggcaaacc atggcccaca ggccaaatcc agacttcacc 1740
tattttcgta aatgaagttt cattggaatg cacacacacc catttgttta catataatat 1800
atggctctct tcattctaaa atgacagagt tgagttagtg caacagagac cccaccgttt 1860
gcaaatcata acatatttac tattttgccc ccttcagaaa gctttccaaa tcctgggtgc 1920
tggcgatttc caatagtctt ctcccgatc ttctgggtta catttctccc tgaaccatc 1980
ctcccatcc ctaaaatttc ccccaaatgt gaactcagtc ataccagttt gctccttatg 2040
tttcattggc ccttgctgct aagagcatcc gcttgacct tctgtctatc cccagacaag 2100
ctttgtcctg tgaccataat gaactcttca tgccgttccc aactttagcc catgttatc 2160
ttctgtctg aatatccacc ctcttctctg ttctcaataa taagttcagg ctcttctgt 2220
tctgagaagc cctttctgac ttccacaggc tgaaccactg gcttctgtc ctctacataa 2280
tacttcaatt ccagcattga tctcactcta tcatgatcat gggtttagct gtctgtccct 2340
gccactgtg tgtgttctc ttgagggcag gaacatttgt tttcacttt ttaaaaaacc 2400
tctgttgccc agtctggcat taggaagtgc ccattagggt gttattgctt gttggcgctt 2460
gagctggggc ttgaaggttt ctataatgtg tagcagtga tagaaaacag gcaggtcaga 2520
aaaggtctct gtgcatcaca ccaacatggc acatgtatac atatgtaaca aatctgcatg 2580
ttgtgcacat gtaccctaaa acttaaaagta taataataa aaaattttta aaaaaaaaag 2640
aagaggtctc ctggaggaga tgacagctga gctaagtcct ggaggatgag aaggagtata 2700
aaataagata ataggagaaa aaaggcagta ggaacagcat gggtaaaagg gatgaggcct 2760
gaaagaggca cgtggaagga aagacaaatg caggaagggg gaatgggagg gaatgctggg 2820
gtacaggcca aagagggagg cattcgggga gtgtctctt aaacatctt tgggtgtgta 2880
atgtgtggtc attgggaaac catggaccgt ttttttttg caaacctcat atccctgtt 2940
acaactgtct gtttgcatgt cagccttga aaagccctt aaggtatcaa ctatgtttt 3000
gttttgcatt cattcaatcc taagtgcaca gaattccggg catattacag gttcccatg 3060
aatgtttctt tctttattaa aatgtatgaa aactctccag atttaaggaa ggtcctcaat 3120
gtttcaaat ctttttgtta gatcattggt cctgtctaca gctgtcaca atttaaggac 3180
tctggttata tttaatcttc acttttgaat tttctgctg aaaaatttgt attagaaaaa 3240
aaagtctatc cttttatgga cggctctaact ctcttgaatc atttgggtt gcttttctt 3300
ggaccttctt caactctgtt ttgtctctgt tgagttaagg cttttaagaa cacctgaatt 3360
ctttctctt gcaaaaccag aggcagctt tttccgcct attttcagtt tatttcttgt 3420
gatttttagt tttttctctt aaccaaagtc taaatggatt taggagaaat aaacttattt 3480
gtaaagctgt caagggacca ttagaaggat ggtgcttcac agatagaata cagtttttat 3540
taatgatgcc tagacaaatc ctgccattag cccaagggt cagaaaagta gcagcctagt 3600
agttttgag ttgtcaatga aatgaattgg actggatggt taaggatgcc cagaagattg 3660
aataaaattg ggatttagga ggaccttgt actccaggaa attctccaag tctccactta 3720
gttatccaga tctcaaatg gaacatgaag cttcagttt aaattgaata cattttccat 3780
ccatggattg gcttgtttt ttcagttgag tgcttgagg tgtctttct acgtaacagc 3840
taaacccacg gcttctcttc tegtaaaacc aaaacaaaaa ggtttctat tcaagtgcct 3900
tctgtgtgtg cacatgtgta atacatatct gggatcaaa ctatctatat aaagtcctt 3960
attctgtgtg ggttcaaaca catttcaaag cttcaggatc ctgaaagggt ttgtctact 4020
tcctgaagac ctgaacaccg ctccataaaa gccatggctt gccttggatt tcagcggcac 4080
aaggctcagc tgaacctggc taccaggacc tggccctgca ctctctgtt tttctcttc 4140

ttccatccctg tcttctgcaa aggtgagtg gacttttggg gcatgaagat ggaggaggtg 4200
ttcttcttac ctgggtttca tttgtttcag cagtcaaaag cagtjattta tagcaaaagcc 4260
agaagttaaa ggtaaaaact caatctggct tggctggctc tgtattccag ggccagcagg 4320
gagcagttgg gcggcacgaa ataaggcaaa gagatagctc agaacagagc gccagggtatt 4380
tagtaggggc ttcattgaatg catgtgagct ggtttagtag agagacacag gcaaattcag 4440
accttcttat gagactggaa gtgatttaag agggaaagga tagccatagt cctgaataca 4500
tttgagctgg gtttcaggat gagctcacia gttcttttaa aaaaaattga cttaaagcaaa 4560
tcttggaag agtttttttg ctatacaatt caagggtttta aggtcctcgg attcatatac 4620
tttataaatg aattagccag cttgtttaaa atgtagggaa attgtgggaa gaatgccttc 4680
tttacttaac tcaagggtttt aagggtctct taatcaattc tactagctaa tttagccaatt 4740
atttaaaaaat aaaagtttga aattgccaaa aaaaaagac aaggaaaagg aaagaaagaa 4800
agccaccagt ctgtttggca tacaatactt aattgttgcc tgacctacgt gtgggtttca 4860
gatgcagatc ctgagttttc agctcttcag agactgacac caggtttgtt acacggctta 4920
aaatgatgag tatatccatt gaatctcaac cttatctctc tctagacctt cttggttaag 4980
aaaccatgta gtttgtatga agtaggtact caaaagatat ttgatgattt aatttttact 5040
ggagaagaaa tattcatata tgtttttctta tttttacatg ttttaaatat gtaaagatta 5100
aataaacact cttagaagta tttaaatttc ctaaagtaaa tttatctcaa ccagtaacag 5160
gacctccca atactggaaa gttgagtggt accgcattta gtggtgatga gtgtgagctt 5220
gcttggggag agggcaggac atttaggatt tcttaagctt agagtcaata caataaagat 5280
tattgagtg cacttgggt gggctataat cactgctcac aggagttcat gaaccacaag 5340
taaaagagtg aggagatatg attagctcac aaataacttt aatacagagc agaaagtaat 5400
gaactactgc aatggagtta tcacagtgtt aaggatgttc agagggcatc tctgataggc 5460
agaggtgagg gtttagggaa gaagctgtag tctagctagc tagagctgct ggaatagaca 5520
tgacaatggc tgctgcaaac tgttttctct tctgaggaca gatgtcccgt gcaagtggct 5580
tggtggaagg gactagtgtc tctaataatg ggtgatttat aagcaggaaa gtgtgtccta 5640
gaaattcaga ccagagtgat agattggaat tggatcatgg gggactcat gaattgtatt 5700
tattgtattt gttttttcga tcagtgttag taaagtgtca aagggttga gcagatgagt 5760
gacatcatgc aacacaagtt ttgagtttca cttgtcagac tgactggaga ggggcctggt 5820
tagttacagg aaggtaattt ggcatgcagc cactattttt gagttgatgc aagcctctct 5880
gtatggagag ctggtctcct ttatctctgt ggaaaagaga acaaaggagc atgggagtg 5940
tcaagggaag gagaataaaa gggcagagag gcagcgggtg tgcagggga agcccacagg 6000
agttaacagc aggggtgcct caacctagag aggaagcgac ctggtgccct cggtctctgt 6060
gcttcttca tctaacaaca tcttccactc tacaacaatg ccagggaagg cgagggtg 6120
tacagtgcac caagacacag ctactcctgg gtgacagagg ttccaggcca gctcactaag 6180
taggcagaag tttttgacat atactttgag agataaagca agattctgt cctcaacctt 6240
cagaatttcc cctaccactc attatagtgc cggagctata tagctcctat cattctatca 6300
taaccttaga ataccagaga acatatcatc tcatttaatt atctcttact atatgtgaaa 6360
aaaatgaagg acatggggga agtggtgactt gccccaaatc acatatttca tggtagagcc 6420
aggtcttctg tttgtcatat cagtgttctt cctgccaaa ccatcttgaa gaatctattt 6480
ctcagtaaga aaatatcttt atggagagta gctggaaaac agttgagaga tggaggggag 6540
gctgggggtg tggagagggg aagggttaag tgatagattc gttgaagggg ggagaaaagg 6600
ccgtggggat gaagctagaa ggcagaaggg cttgcctggg cttggccatg aaggagcatg 6660
agttcactga gttccctttg gcttttccat gctagcaatg cagtgggccc agcctgctgt 6720
ggtactggcc agcagccgag gcatcgccag ctttgtgtgt gagtatgcat ctccaggcaa 6780
agccactgag gtccgggtga cagtgtctcg gcaggctgac agccagggtga ctgaagtctg 6840
tgccgcaacc tacatgatgg ggaatgagtt gaccttcta gatgattcca tctgcacggg 6900
cacctccagt ggaaatcaag tgaacctcac tatccaagga ctgagggcca tggacacggg 6960
actctacatc tgcaagggtg agctcatgta cccaccgcca tactacctgg gcataggcaa 7020

cggaacccag atttatgtaa ttggtagca aagccatttc actgagttga caccctgttc 7080
 attgcagctt tctatgcaca aaaacagttt tgttccctaa ttcaggagg tttaccttta 7140
 ggactgtgga catttctctt aagagttctg taccacatgg tagccttgc tttgtgggt 7200
 ggcaacctta atagcattct gactgtaaaa taaaatgat tggggaagt ggggtctctg 7260
 ctctggagtg ctaaccatca tgacgttga tctgtacctt tgatatgata tgatgctctt 7320
 ggggaagttag tcccaaatag ccaaacctat tgggtgggcta cccatgcaat ttagggttg 7380
 acctcaaggc ctggaagctc taatgtctct ttttcccaa tgttggggag tagagcccta 7440
 gagttaaaaa ctgtctcagg gaggtctctg tttgtttctt gttgcagatc cagaacctgt 7500
 cccagattct gacttctctc tctggatctt tgcagcagtt agttcgggtt tgttttttta 7560
 tagctttctc ctacagctg tttctttgag caaaatgggt agtgtgtgtc tgatggtgca 7620
 ccatgtctga tggggatacc tttagtgtta tcaactggcc aaaagatgat gttgagttta 7680
 gtgttcttga gatgagatga ggcaataaat gaagaggag gacagtgtta aagaacgcac 7740
 tagaaccgta ggcatgggca tttgaggttt cagaatgact aatattttag atgaatttgt 7800
 ttgacattga atgttcatgt gcttctgagc aggggttcaa tttgagtaac cgttgcaata 7860
 acatggggca gctgttttgc tctttgtctt catgacaact gtacttaagc taacagccct 7920
 gaaacatgag attaggctgg gcagaatgct gctagagagg accacttga tggctcttat 7980
 tctctctctc catgtccctc tccatcacct ggaagtcacc tctgggtgcc actctggtgc 8040
 ctctcttctc gaagctgtag ctgtctcatc gacacctatc cctgttatcc agtttgcctg 8100
 actgggagct tttgccttcc ccttcagcca ggaagtgaat gtcccagttt ttatttatca 8160
 caggtgttgg tactgtgtgt agaagaggta gaattatgga atcaggccctc ctgtcaggat 8220
 tttcttttga cagtccctct cagacacctc tgcctaaggc cagctttgcc attacaaact 8280
 ctccctctc cctctctctc tctctctctt cctcttctct ctctctctc tttctctctc 8340
 tctcttctc cctctctgtc tcttatacac atacacaaag atatactcta tccaacatc 8400
 ctctacccaa cctgacagag atgtctcttg ctgtaggttc agcagtggg atgagaaata 8460
 cagctctcaa acaggataac taaagcttat tatcttatca agcttgttcc cttgcagaca 8520
 agattgatca attatcatag gctttctggg tgttctttct gaagctttct caaagtctct 8580
 tttctctatc ttccattcaa ggcaaatgat tgccatttaa catcaaaatc acagtatttt 8640
 atctaaaata aattttaata gctgaatcaa gaaaatctcc tgaggtttat aattctgtat 8700
 gctgtgaaca ttcattttta accagctagg gacccaatat gtgttgagtt ctattatggt 8760
 tagaagtggc ttccgtatct ctcatagta attactgtt ctttttgtgt ttgacagcta 8820
 aagaaaagaa gccctcttac aacaggggtc tatgtgaaaa tgccccaac agagccagaa 8880
 tgtgaaaagc aatttcagcc ttattttatt cccatcaatt gagaaacct tatgaaag 8940
 agagtccata tttcaatttc caagagctga ggcaattcta acttttttgc tatccagcta 9000
 ttttattttg tttgtgcatt tggggggaat tcatctctct ttaataataa gttggatgcg 9060
 gaacccaaat tacgtgtact acaatttaaa gcaaaggagt agaaagacag agctgggatg 9120
 tttctgtcac atcagctcca ctttcagtga aagcatcact tgggattaat atggggatgc 9180
 agcattatga tgtgggtcaa ggaattaagt taggggaatgg cacagcccaa agaaggaaag 9240
 gcaggagcg agggagaaga ctatattgta cacaccttat atttacgtat gagacgttta 9300
 tagccgaat gatctgttca agttagatgt tatgcc 9336

<210> 2

<211> 672

<212> DNA

<213> Homo sapien

<400> 2

atggcttgc tttgatttca gcggcacaag gctcagctga acctggctac caggacctgg 60

cccatgcaactc tccctgttttt tcttctcttc atccctgtct tctgcaaagc aatgcacgtg 120
 gcccagcctg ctgtggactt ggccagcagc cgaggcatcg ccagctttgt gtgtgagtat 180
 gcatctccag gcaaagccac tgaggcccg gtgacagtc ttcggcaggc tgacagccag 240
 gtgactgaag tctgtgcggc aacctacatg atggggaatg agttgacctt cctagatgat 300
 tccatctgca cgggcacctc cagtggaaat caagtgaacc tcaactatcca aggactgagg 360
 gccatggaca cgggactcta catctgcaag gtggagctca tgtaccacc gccatactac 420
 ctgggcatag gcaacggaac ccagatttat gtaattgac cagaaccgtg cccagattct 480
 gacttctctc tctggatcct tgcagcagtt agttcggggt tgttttttta tagctttctc 540
 ctcacagctg tttctttgag caaaatgcta aagaaaagaa gccctcttac aacaggggtc 600
 tatgtgaaaa tgcctccaac agagccagaa tgtgaaaagc aatttcagcc ttattttatt 660
 cccatcaatt ga 672

<210> 3

<211> 223

<212> PRT

<213> Homo sapien

<400> 3

Met Ala Cys Leu Gly Phe Gln Arg His Lys Ala Gln Leu Asn Leu Ala
 1 5 10 15

Thr Arg Thr Trp Pro Cys Thr Leu Leu Phe Phe Leu Leu Phe Ile Pro
 20 25 30

Val Phe Cys Lys Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala
 35 40 45

Ser Ser Arg Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly
 50 55 60

Lys Ala Thr Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln
 65 70 75 80

Val Thr Glu Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr
 85 90 95

Phe Leu Asp Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val
 100 105 110

Asn Leu Thr Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile
 115 120 125

Cys Lys Val Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly
 130 135 140

Asn Gly Thr Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser
 145 150 155 160

Asp Phe Leu Leu Trp Ile Leu Ala Ala Val Ser Ser Gly Leu Phe Phe
165 170 175

Tyr Ser Phe Leu Leu Thr Ala Val Ser Leu Ser Lys Met Leu Lys Lys
180 185 190

Arg Ser Pro Leu Thr Thr Gly Val Tyr Val Lys Met Pro Pro Thr Glu
195 200 205

Pro Glu Cys Glu Lys Gln Phe Gln Pro Tyr Phe Ile Pro Ile Asn
210 215 220

<210> 4
<211> 15
<212> DNA
<213> Homo sapien

<400> 4
agatcctyaa agtga

15

<210> 5
<211> 15
<212> DNA
<213> Homo sapien

<400> 5
cagtcaargg cagtg

15

<210> 6
<211> 15
<212> DNA
<213> Homo sapien

<400> 6
cactgagytg acacc

15

<210> 7
<211> 15
<212> DNA
<213> Homo sapien

<400> 7
ctagaacygt aggca

15

<211> 15
<212> DNA
<213> Homo sapien

<400> 13
ataaatcact gccyt

15

<210> 14
<211> 15
<212> DNA
<213> Homo sapien

<400> 14
ccatttcact gagyt

15

<210> 15
<211> 15
<212> DNA
<213> Homo sapien

<400> 15
gcaacaggtg tcarc

15

<210> 16
<211> 15
<212> DNA
<213> Homo sapien

<400> 16
aacgcactag aacyg

15

<210> 17
<211> 15
<212> DNA
<213> Homo sapien

<400> 17
tgccaatgcc tacrg

15

<210> 18
<211> 15
<212> DNA
<213> Homo sapien

<400> 18

aataaatttt aatrg

15

<210> 19

<211> 15

<212> DNA

<213> Homo sapien

<400> 19

ttcttgattc agcya

15

<210> 20

<211> 15

<212> DNA

<213> Homo sapien

<400> 20

ctgtatgctg tgarc

15

<210> 21

<211> 15

<212> DNA

<213> Homo sapien

<400> 21

ttaaaaatga atgyt

15

<210> 22

<211> 10

<212> DNA

<213> Homo sapien

<400> 22

tccagatcct

10

<210> 23

<211> 10

<212> DNA

<213> Homo sapien

<400> 23

tgttcacttt

10

<210> 24
<211> 10
<212> DNA
<213> Homo sapien

<400> 24
cagcagtcaa

10

<210> 25
<211> 10
<212> DNA
<213> Homo sapien

<400> 25
aatcactgcc

10

<210> 26
<211> 10
<212> DNA
<213> Homo sapien

<400> 26
tttcactgag

10

<210> 27
<211> 10
<212> DNA
<213> Homo sapien

<400> 27
acaggtgtca

10

<210> 28
<211> 10
<212> DNA
<213> Homo sapien

<400> 28
gcactagaac

10

<210> 29

<211> 10
<212> DNA
<213> Homo sapien

<400> 29
caatgcctac

10

<210> 30
<211> 10
<212> DNA
<213> Homo sapien

<400> 30
aaattttaat

10

<210> 31
<211> 10
<212> DNA
<213> Homo sapien

<400> 31
ttgattcagc

10

<210> 32
<211> 10
<212> DNA
<213> Homo sapien

<400> 32
tatgctgtga

10

<210> 33
<211> 10
<212> DNA
<213> Homo sapien

<400> 33
aaaatgaatg

10

<210> 34
<211> 18
<212> DNA
<213> Homo sapien

<400> 34
tgtaaaacga cggccagt

18

<210> 35
<211> 19
<212> DNA
<213> Homo sapien

<400> 35
aggaaacagc tatgaccat

19

<210> 36
<211> 9336
<212> DNA
<213> Homo sapien

<220>
<221> allele
<222> (3735)
<223> PS1: polymorphic base C or T

<220>
<221> allele
<222> (4102)
<223> PS2: polymorphic base A or G

<220>
<221> allele
<222> (4238)
<223> PS3: polymorphic base A or G

<220>
<221> allele
<222> (7067)
<223> PS4: polymorphic base T or C

<220>
<221> allele
<222> (7747)
<223> PS5: polymorphic base C or T

<220>
<221> allele
<222> (8660)
<223> PS6: polymorphic base A or G

<220>

<221> allele

<222> (8708)

<223> PS7: polymorphic base A or G

<400> 36

```

actcaaatcc cccccagttt ctcagtaatc tgctttgtat ttttttttc aatcaataat 60
aacacattaa attctgtcat catgtctcta tttctcctt taatctagaa tgggtttcca 120
ggcttttccc tttttaatct tttgtgatg taacattttt gaagaatcta ggccagttat 180
tttgagaatc atggatttct ctgatttttt cttcatgctt agattcagtt aaaacaattt 240
tggcaataac tacacaaagt tgttgtgtcc ttctcagtg attgtatctg gaggtacctt 300
atgtcagaac actccatcac agtaatcttc agttgggac acttggttaa agcagtatct 360
gctagattta tcaattatct tttccctatt gtaattatca agtaatctgt ggggtggtat 420
atgtgaatat atcttttctc accaagattc ttttaccag catttttttt acaaccattt 480
gaattttctt gctatgacag aattagtttc ataagagcat aattaaggaa tctggttaaga 540
cgtctttgaa agttctaat acaacctca tctctcttta tgttaaactg tcttcattca 600
tttacctacc ttttaacaa atgtttaaag aatgctcgtt ttgtggcttg gcctgtgaca 660
ttttcatcag tggcttcata gttgcctaaa ctaatgtctc ttctgtgct ccagaatcct 720
attggcaact attttccatg gatgccataa actcaacata ctcagatgaa agctgtcatt 780
tcactttctc acctgctctt cctgttattt attccagtaa attgcacctc tgtcttctca 840
ttgaccttat aataagaaag gcctatttat tgagtggcta tctgttacia taggcctttt 900
atacatgtta catgcattat ctcatttaat cttttcaata atcttgagaa ttagatgcca 960
ttatctggag gacattgggc tttgagcccc aggtcaacag taagcaagtc cccatgctag 1020
gatacaaatc cagttatctt gggcttcaaa atctttgctt tagcatagtg ataaaggat 1080
cactggcttc aaatcctggc tctgccactt cctacgggtg attgggtaag tcaccttaac 1140
tttcaatgcc ttgatttctt tctttataaa atgggaaaaa tggtaactct tgtctttag 1200
ggttgttatg gacttgaaag aaacaaaact tgctgggccc ggtggctcac gcctgtaac 1260
ccagcgcttt gggaggccga ggcgggtaga tcacctgagg tcaggaattt gagactagcc 1320
tggcacaacat ggcgaaaccc catctctact aaaataatac aaaaaattag ctgggctggt 1380
tgggtgggtgc ctgcaatccc agctacttgg gaggtgagg caggagaatc acttgaaccc 1440
aggaggcaga ggttgacgtg agccgagatc gcgccactgc actccaacct gggaaacaa 1500
agtgaactc tgtctcaaaa aaataaaaaa gaaaagaaag aaaacttgac atttgaataa 1560
tttctgcagc accacagtac ttctcaaac ataattctgg tgtcattcat tctaaacttc 1620
tccctccact cactgattgt ggactagtcc catagggacc tctcttcac agctgtcatt 1680
accttagttt agaccagggg ttggcaaac atggcccaca ggccaaatcc agacttcacc 1740
tattttctga aatgaagttt tattggaatg cacacacacc catttggtta catataatat 1800
atggcttctt tcattctaaa atgacagagt tgagttagtg caacagagac cccaccgttt 1860
gcaaatcata acatatttac tattttgccc ctttcagaaa gctttccaaa tcttgggtgc 1920
tggcgctatt caatagtctt cttcccagtc ttctgggtta catttctccc tgaacccatc 1980
ctcccatcc ctaaaatttc ccccaaatgt gaactcagtc ataccagttt gctccttatg 2040
tttcattggc ccttgcctgt aagagcatcc gcttgacact tctgctcatc cccagacaag 2100
ctttgtctgt tgaccataat gaactcttca tgccgtttcc aacttttagcc catgttatcc 2160
ttctgtctgt aatatccacc cttttctctg ttctcaataa taagtccagg cttttctgtc 2220
tctgagaagc cttttctgac ttccacagcg tgaacctg gcttctgctc ctctacataa 2280
taactcaatt ccagcattga tctcactcta tcatgatcat gggtttagct gtctgtccct 2340
gccactgctg tgtgttcttc ttgagggcag gaacatttgt ttttcacttt ttaaaaaacc 2400
tctgttgccc agcttgccat taggaagtgc ccattaggtt gttattgctt gttggcgctt 2460

```

gagctggggc ttgaagggtt ctataatgt tagcagtgta tagaaaacag gcaggtcaga 2520
 aaaggcttct gtgcattcaca ccaacatggc acatgtatc atagttaaca atcttgcattg 2580
 ttgtgcacat gtacctaaa acttaagta taataaata aaaattttaa aaaaaaaaag 2640
 aagagggctt ctggaggaga tgacagctga gctaagctt ggaggatgag aaggagtata 2700
 aaataagata ataggagaaa aaaggcagta ggaacagcat gggtaagggt gatgaggcct 2760
 gaaagaggca cgtggaagga aagacaaatg cagggaaggg gaatgggagg gaatgctggg 2820
 gtacaggcca aagagggagg cattcgggga gtgttctt aaacatcctt tgggtgtgcta 2880
 atgtgtggtc attgggaaac catggaccgt ttttttttgg caaacctcat atccctgtt 2940
 acaactgtct gtttgcatgt cagccttgta aaagccctt aaggtatcaa ctatgtttt 3000
 gttttgcatt cattcaatcc taagtgcaca gaattccggg catattacag gttcccatg 3060
 aatgtttctt tctttattaa aatgtatgaa aactctccag atttaaggaa ggtcctcaat 3120
 gtttcaaatt ctttttgtta gatcattggt cctgtctaca gctgtcacia atttaaggac 3180
 tctggttata tttaatcttc acttttgaat tttctgctt gaaaatttgt attagaaaaa 3240
 aaagtctatc cttttatgga cggctcctat ctcttgaatc atttgggttg gcttttctt 3300
 ggacctctt caactctgtt ttgtctctgt tgagttaagg cttttaagaa cacctgaatt 3360
 cttctcttct gcaaaaccag aggcagcttc ttttccgct attttcagtt tatttcttgt 3420
 gattttagtt tttttctctt aaccaaatgc taaatggatt taggagaaat aaacttattt 3480
 gtaaagctgt caagggacca ttagaaggat ggtgcttcac agatagaata cagtttttat 3540
 taatgatgcc tagacaaatc ctgccattag ccaagggtc cagaaagtta gcagcctagt 3600
 agttttggag ttgtcaatga aatgaattgg actggatggg taaggatgcc cagaagattg 3660
 aataaaattg ggatttagga ggaccttgt actccaggaa attctccaag tctccactta 3720
 gttatccaga tccctyaaagt gaacatgaag cttcagtttc aaattgaata ctttttccat 3780
 ccatggattg gcttgttttg ttcagttgag tgcttgaggt tgtcttttc acgtaacagc 3840
 taaaccacag gcttcttctc tctgtaaaacc aaaaacaaaa ggctttctat tcaagtgcct 3900
 tctgtgtgtg cacatgtgtg atacatatct gggatcaaag ctatctatat aaagtccttg 3960
 attctgtgtg ggttcaaaca ctttcaaag cttcaggatc ctgaaagggt ttgctctact 4020
 tctgaagac ctgaacaccg ctcccataaa gccatggctt gccttggtt tcagcggcac 4080
 aaggctcagc tgaacctggc tccaggacc tggccctgca ctctctgtt ttttcttctc 4140
 ttcattccctg tcttctgcaa aggtgagtga gacttttga gcatgaagat ggaggagggtg 4200
 tttctctac ctgggtttca tttgtttcag cagtcaargg cagtgattta tagcaaagcc 4260
 agaagttaa ggtaaaactc caatctggct tggctggctc tgtattccag ggccagcagg 4320
 gagcagttgg gcggcacgaa ataaggcaa gagatagctc agaacagagc gccagggtatt 4380
 tagtaggggc ttcattgaatg catgtgagtt ggtttagtag agagacacag gcaatttcag 4440
 acccttctat gagactggaa gtgatttaag agggaaagga tagccatagt cctgaatata 4500
 tttgagctgg gtttcaggat gagctcacia gttcctttaa aaaaaattga ctttaagcaa 4560
 tcttggaag agtttttttg ctatacaatt caaggtttta aggtcctcgg attcatatac 4620
 tttataaatg aattagccag cttgttttaa atgtaggga attgtggga gaatgccttc 4680
 tttacttaat tcaaggtttt aaggttctct taatcaattc tactagctaa tttagccaatt 4740
 atttaaaaat aaaagtttga aattgccaaa aaaaaagac aaggaaaagg aaagaaagaa 4800
 agccaccagt ctgtttggca tacaatactt aattgttgcc tgacctacgt gtgggtttca 4860
 gatgcagatc ctcagttttc agctcttcag agactgacac caggtttgtt acacggctta 4920
 aaatgatgag tatatccatt gaatctcaac cttatctctc tctagacctt cttgggttaag 4980
 aaacctatga gtttgtatga agtaggtact caaaagatat ttgatgattt aatttttact 5040
 ggagaagaaa tattcatata tgttttctta tttttacatg ttttaaatat gtaaagatta 5100
 aataaacact cttagaagta tttaaatctc ctaaaagtaa tttatctcaa ccagtaacag 5160
 gacctccca atactggaaa gttgagtggt accgcattta gtgggtatga gtgtgagctt 5220
 gcttggggag agggcaggac atttaggatt tcttaagctt agagtcaata caataaagat 5280
 tattgagtg cactctgggt gggctataat cactgctcac aggagttcat gaaccacaag 5340

taaaagagtg aggagatatg attagctcac aaataacttt aatacagagc agaaagtaat 5400
 gaactactgc aatggaggtta tcacagtgc aaggatgctc agagggcatc tctgataaggc 5460
 agaggtgagg gttagggaag gaagctgtag tctagctagc tagagctgct ggaatagaca 5520
 tgacaatggc tgcctgcaaac tgttttctct tctgaggaca gatgtccctt gcaagtggct 5580
 tgggtggaagg gactagtgtc tctaatatag ggtgatttat aagcaggaaa gtgtgtccta 5640
 gaaattcaga ccagagtgat agactggaat tggatcatgg gggactcatt gaattgttatt 5700
 tatctgattt gtcttttgcga tcagtgttag taaagtgtca aagggtattga gcagatgagt 5760
 gacatcatgc aacacaagtt ttgagtttca cttgtcagac tgactggaga ggggcttgg 5820
 tagttacagg aaggtaatct ggcatgcagc cactattttt gagtgtatgc aagcctctct 5880
 gtatggagag ctggtctcct ttatcctgtg ggaanaagaga acaaaggagc atgggagtg 5940
 tcaagggaag gagaaataaa gggcagagag gcagcgttg tgtcagggga agcccacagg 6000
 agttaacagc aggtttgctt caacctagag aggaagcgac ctggtgccct cggctctgtg 6060
 gcttcttcca tctaacaaca tcttccactc tacaacaatg ccagggaagg cggaggcttg 6120
 tacagtgcac caagacacag ctactcctgg gtgacagagg ttcagggccca gctcactaag 6180
 taggcagaag tttttgacat atactttgag agataaagca agattctgta cctcaacctt 6240
 cagaatttcc cctaccactc attatagttc cggagctata tagctcctat cattctatca 6300
 taaccttaga ataccagaga acatatcatc tcatctaatt atctcttact atatgtgaaa 6360
 aaaatgaagg acatggggga agtgtgactt gccccaaatc acatatttca tggtagagcc 6420
 aggtcttctg tttgtcatat cagtgttctt cctgccacaa ccatcttgaa gaattctatt 6480
 ctcagtgaag aaatatcttt atggagagta gctggaaaac agttgagaga tggaggggag 6540
 gctgggggtg tggagagggg aagggttaag tgatagattc gttgaagggg ggagaaaagg 6600
 ccgtggggat gaagctagaa ggcagaaggg cttgcctggg cttggccatg aaggagcatg 6660
 agttcactga gttccctttg gcttttccat gctagcaatg cacgtggccc agcctgctgt 6720
 ggtactggcc agcagccgag gcatcgccag ctttgtgtgt gagtatgcat ctccaggcaa 6780
 agccactgag gtccgggtga cagtgtctcg gcaggctgac agccagggtga ctgaagtctg 6840
 tgccgcaacc tacatgatgg ggaatgagtt gaccttccca gatgattcca tctgcacggg 6900
 cacctccagt ggaaatcaag tgaacctcac tatccaagga ctgagggcca tggacacggg 6960
 actctacatc tgcaagggtg agctcatgta cccaccgcca tactacctgg gcataggcaa 7020
 cggaacccag atttatgtaa ttggtgagca aagccatttc actgagytga cacctgttgc 7080
 attgcagtct tctatgcaca aaaacagttt tgttccctaa tttcaggagg tttactttta 7140
 ggactgtgga cattctcttt aagagtcttg taccacatgg tagccttgct tattgtgggt 7200
 ggcaacctta atagcattct gactgtaaaa taaaatgatt tggggaagtt ggggctctcg 7260
 ctctggagtg ctaacctca tgacgtttga tctgtacttt tgatatgata tgatgtcctt 7320
 ggggaagtag tcccaaatag ccaaacctat tgggtgggcta cccatgcaat ttaggggttg 7380
 acctcaaggc ctggaagctc taatgtcctt ttttcaccaa tgttggggag tagagcccta 7440
 gagttaaaa ctgtctcagg gaggtctctc tttgtttctt gttgcagatc cagaacctg 7500
 ccagattct gacttctctc tctggatcct tgcagcagtt agttcggggg tgttttttta 7560
 tagctttctc ctcacagctg tttctttgag caaatgggtg agtgtgtgtc tgatggtgca 7620
 ccatgtctga tggggatacc tttagtggta tcaactggcc aaaagatgat gttgagttta 7680
 gtgttcttga gatgagatga ggcaataaat gaagaggaag gacagtggta aagaacgcac 7740
 tagaacygta ggcattggca tttgaggttt cagaatgact aatattttag atgaatttgt 7800
 ttgacattga atgttcatgt gcttctgagc agggtttcaa tttgagtaac cgttgcaata 7860
 acatggggca gctgttttgc tctttgtctt catgacaact gtacttaagc taacagccct 7920
 gaaacatgag attaggctgg gcagaatgct gctagagagg accacttgga tggctctttat 7980
 tctcttctc catgtccctc tccatcacct ggaagtcacc tctgggtgcc actctggtgc 8040
 cttcttctc gaagctgtag ctgctcacat gacacctatc cctgttatcc agtttgcctt 8100
 actgggacgt tttgccttcc ccttcagcca ggaagtgaat gtcccagttt ttatttatca 8160
 cagggtgttg tattgggtgt agaagaggta gaattatgga atcaggccctc ctgtcaggat 8220